



History: 8 year old girl with history of limping.



Magnified AP view of the right hip






Coronal IR image of both hips

DEPARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY



Coronal PD images through the right hip

HOSPITAL FOR SPECIAL SURGERY DEPT
DEPARTMENT OF
DEPARTMENT OF
DEPARTMENT OF



Coronal SPGR (physal sensitive sequence) images of the right hip

Findings

Radiographs of the pelvis demonstrate coxa vara of the right hip. An irregular, widened right capital femoral physis is present with a vertical orientation and Y shaped configuration of the physis. There is also a triangular focus of ossification along the inferomedial aspect of the physis. Scanogram shows a 2.5cm limb length discrepancy. MR images demonstrate a widened, irregular, proximal femoral physis with the inferomedial, triangular focus of ossification. A mild degree of edema is seen about the right, proximal femoral physis.



Widened, irregular right capital femoral physis with vertical orientation and Y shaped configuration.

Normal physis




Widened, irregular proximal femoral physis

Triangular ossification at inferomedial aspect of the physis



Thickened, irregular physis



Y shaped configuration of physis

Triangular focus of ossification

DEPARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY DEPARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY DEPARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY DEPARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY DEPARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY



Irregular, widened femoral physis




Triangular focus of ossification

Diagnosis: Infantile or congenital coxa vara (CCV)

Infantile or congenital coxa vara results from abnormal maturation of the proximal femoral physis that causes decreased ossification, weakening of the bone, and subsequent coxa vara. Classically, as in this case, there is a Y shaped configuration of the proximal femoral physis and a focus of ossification inferomedially. Patients typically present from the time they start walking up to about 6 years of age and present with a limp.

There is often a mild limb length discrepancy of 2-4 cm as is seen in this case. If necessary, a valgus osteotomy is performed. Osteotomy is indicated for a Hillgenreiner epiphyseal angle (HEA) of > 60 degrees or an HEA of 45-60 degrees with increasing coxa vara. In distinction, to proximal femoral focal dysplasia (PFFD), the varus deformity in CCV is at the level of the physis and not subtrochanteric as in PFFD. Also, in PFFD, there is typically a more pronounced limb length discrepancy.

HEA: Angle subtended by a line between the triradiate cartilages (1) and a line of the proximal femoral physis (2). Notice the very high (>60 deg) angle on the right and the normal angle on the left of 20-25 degrees.



Postoperative



Marked reconstituion of alignment

Preoperative



Postoperative



Resources

- <http://www.posna.org/education/StudyGuide/coxaVara.asp>
- <http://emedicine.medscape.com/article/1259556-overview>
- **Very special thanks to Roger Widmann, MD for his assistance in this case presentation.**

