






Sagittal PD of the involved finger


History: 42 year old with three years of finger pain



Axial IR



Axial PD



## Findings

A high signal mass is present about the dorsum of the distal phalanx where it has caused indolent erosion along the dorsum of the distal phalanx. The mass demonstrates marked high signal on the inversion recovery sequence and high signal on the PD sequences. Although, not employed in this case, the mass would typically demonstrate marked enhancement following gadolinium administration.




D IMAGING HOSPITAL FOR  
RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY DEPARTMENT OF  
ARTMENT OF RADIOLOGY AND IMAGING HOSPITAL FOR SPECIAL SURGERY DEPARTMENT  
AND IMAGING HOSPITAL FOR SPECIAL SURGERY



High signal mass

Indolent erosion of the dorsum of the distal phalanx



High signal mass

Erosion of distal phalanx



## Diagnosis: Glomus Tumor

- A glomus tumor is a hamartoma (tissue normally found at a location but growing in a disorganized pattern) of the neuromyoarterial (nm) apparatus. Typically found in women from 30 to 50 years of age, although small, the masses are intermittently, extremely painful. Patients have often 2-3 years of symptoms before seeking treatment. They are as in this case, high in signal and often yield erosion of the underlying bone. Given the classic appearance in this case, no contrast was administered, but as the mass represents proliferation of the nm apparatus, it will avidly enhance. They typically occur at the subungual location as in this case but occasionally extend into the underlying bone.



## References:

- [http://www.wheelessonline.com/ortho/glomus\\_tumor](http://www.wheelessonline.com/ortho/glomus_tumor)
- <http://www.bonetumor.org/tumors-vascular-tissue/glomus-tumor>

