

## **SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES**

### **(Anterior Capsule Repair, Bankart Repair)**

The following post-operative shoulder anterior stabilization guidelines were developed by HSS Rehabilitation and are categorized into five phases with the ultimate goal for returning the athlete to full competition. They can be used for patients undergoing a variety of anterior stabilization procedures with attention given to exact location of repair and any concomitant procedures. It is important that full range of motion is restored while respecting soft tissue healing. Classification and progression are both criteria-based and time based due to the healing constraints of the human body. The first phase is focused on soft tissue healing and maintenance of pain-free range of motion (ROM). Phases two and three are focused on building foundational strength and stability which will allow the athlete to progress to phase four which includes plyometric exercises. With the completion of phase four the athlete will be able to start the final phase which includes interval sports programs. Cardiovascular endurance, and hip and core strengthening should be addressed through the rehabilitation process. The clinician should use their skilled judgement and decision making as the athlete advances as all progression may not be linear.

### **FOLLOW SURGEON MODIFICATIONS AS PRESCRIBED**

## SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES

### Phase 1: Week 0-1

#### PRECAUTIONS

- Sling for 3 weeks (as per surgeon guidelines)
- Avoid stress on anterior shoulder joint
- If combined with biceps tenodesis, no biceps strengthening for 8 weeks
- No forced passive range of motion (PROM)
- Avoid painful activities

#### ASSESSMENT

- Quick Disabilities of Arm, Shoulder & Hand (Quick DASH)
- Numeric Pain Rating Scale (NPRS)
- Assessment of incision
- Cervical mobility
- Shoulder PROM
- Distal upper extremity PROM
- Palpation
- Static scapular assessment (Kibler Grading)
- Posture assessment

#### TREATMENT RECOMMENDATIONS

- Patient education
- Gripping and hand active range of motion (AROM)
- Postural awareness
- Wrist AROM: flexion/extension/pronation/supination
- ROM
  - Week 1: external rotation (ER) to neutral, elevation in scapular plane 60°

#### CRITERIA FOR ADVANCEMENT

- Decreasing discomfort at rest

#### EMPHASIZE

- Protection of repair
- Reduction of tissue irritability
- Prevention of muscle atrophy

## SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES

### Phase 2: Weeks 2-5

#### PRECAUTIONS

- Sling for 3 weeks (as per surgeon guidelines)
- Monitor for shoulder stiffness
- No forced PROM
- Avoid undue stress to anterior shoulder joint
- If combined with biceps tenodesis, no biceps strengthening for 8 weeks

#### ASSESSMENT

- Quick DASH
- NPRS
- Assessment of incision
- Palpation
- Cervical mobility
- Shoulder PROM
- Static/dynamic scapular assessment (Kibler grading)

#### TREATMENT RECOMMENDATIONS

##### **PROM goals – do not force but assess for stiffness**

- Week 2-3
  - Elevation in scapular plane: 90°
  - ER in scapular plane: 5°-10°
  - Internal rotation (IR) in scapular plane: 30°-45°
- Week 4
  - Elevation in scapular plane: 90°-100°
  - ER in scapular plane: 15°-20°
  - IR in scapular plane: 50°-60°
- Week 5-6
  - Elevation in scapular plane: 120°-145°
  - ER in scapular plane: 40°-60°
  - IR in scapular plane: 50°-60°
- Abduction
  - 0°-90° first 6 weeks (gentle motion)

## **Therapeutic Exercise**

- Week 2
  - Scapular isometrics
  - Elbow AROM
  - Shoulder active assisted ROM
- Week 3
  - Rotator cuff (RC) isometrics
  - Rhythmic stabilization ER/IR with physical therapist
- Week 4
  - Continue RC isometrics
  - Resistance band row, resistance band extension
- Week 5-6
  - RC isotonic if arthroscopic; if open start week 6
  - Scapular strengthening
    - Prone row, prone extension, supine serratus punch

## **CRITERIA FOR ADVANCEMENT**

- No pain at rest
- PROM: 120° shoulder elevation; 45° ER in scapular plane
- Tolerance of scapular and RC exercises without discomfort

## **EMPHASIZE**

- Reduction of tissue irritability
- Activation of RC and scapular stabilizers

## SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES

### Phase 3: Weeks 6-15

#### PRECAUTIONS

- No forced PROM
- Avoid stress to anterior shoulder joint
- No painful activities

#### ASSESSMENT

- Quick DASH
- NPRS
- Cervical mobility
- Thoracic mobility
- Shoulder PROM/AROM
- Palpation
- Static/dynamic scapular assessment (Kibler grading)
- Grip strength: dynamometer

#### TREATMENT RECOMMENDATIONS

##### ROM Goals

- Week 6-7
  - Initiate light and pain free ER at 90° shoulder abduction
    - Progress to 30°
- Week 7-9
  - Shoulder flexion 160°-180°
  - ER at 90° shoulder abduction: 75°-90°
  - IR at 90° shoulder abduction: 70°-75°
- Week 9-12
  - Shoulder flexion 180°
  - ER at 90° shoulder abduction: 100°-115°

##### Flexibility

- Shoulder: posterior capsule stretch at physical therapist discretion

## **Therapeutic Exercises**

- Throwers Ten
- Advanced Throwers Ten
- Scapular stabilization
  - Closed chain quadruped double arm protraction
  - Prone “T, I” and progress to “Y” and “W” as ROM allows
- End range stabilization using exercise blade/perturbations
- Shoulder endurance exercise
- UE ergometry (if ROM allows)
- Core strength/kinetic linking
- Weeks 10-16
  - ER/IR strengthening at 90 degrees of shoulder abduction

## **CRITERIA FOR ADVANCEMENT**

- Full shoulder AROM
- Shoulder manual muscle testing (MMT) 4/5 strength below shoulder height

## **EMPHASIZE**

- Full shoulder PROM and AROM
- Restoration of scapular and RC muscle balance and endurance

## SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES

### Phase 4: Weeks 16-19

#### PRECAUTIONS

- No painful activities
- Full ROM

#### ASSESSMENT

- Quick DASH
- NPRS
- Cervical mobility
- Thoracic mobility
- Shoulder PROM/AROM
- Palpation
- Static/dynamic scapular assessment (Kibler grading)
- Elbow PROM/AROM
- Shoulder manual muscle testing (MMT)
- Grip strength: dynamometer

#### TREATMENT RECOMMENDATIONS

- Continue shoulder RC and scapular stabilization exercises
- Continue and progress all Advanced Thrower's Ten exercises
- Initiate plyometrics as tolerated
  - Plyometric progression (over a 4-week period)
    - Double hand chest pass
    - Double hand overhead soccer pass
    - Double hand chops
    - Single hand IR at 0° shoulder abduction
    - Eccentric catch
    - Single hand 90/90 IR
  - Endurance progression
    - Double hand overhead wall taps
    - Single arm 90/90 wall taps
    - Single arm 12 o'clock to 3 o'clock wall taps
    - Exercise blade at multiple angles

## **CRITERIA FOR ADVANCEMENT**

- Full shoulder AROM
- Shoulder manual muscle testing (MMT) 5/5 strength below shoulder height
- Symptom free progression through plyometrics and endurance program

## **EMPHASIZE**

- Shoulder flexibility, strength and endurance
- Pain free plyometrics
- Kinetic linking



## SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES

### Phase 5: Months 5+ (Return to Sport)

#### PRECAUTIONS

- All progressions should be pain free
- Monitor for loss of strength and flexibility

#### ASSESSMENT

- Quick DASH
- NPRS
- Cervical mobility
- Thoracic mobility
- Shoulder PROM/AROM
- Palpation
- Static/dynamic scapular assessment (Kibler grading)
- Shoulder MMT
- Grip strength: dynamometer

#### TREATMENT RECOMMENDATIONS

- Initiate interval sports programs at 5 months
- Continue with all upper and lower extremity flexibility exercises
- Continue with advanced shoulder and scapular strengthening exercises
- Gradually progress sports activities
- Monitor workload
- Collaborate with ATC, performance coach/strength and conditioning coach, skills coach and/or personal trainer to monitor load and volume with return to sport participation

#### CRITERIA FOR RETURN TO PARTICIPATION

- Symptom free progression through interval sports program
- Independent with all maintenance exercises
- Assess need for HSS Throwing Analysis

#### EMPHASIZE

- Return to sports participation
- Collaboration with Sports Performance experts

## SHOULDER ANTERIOR STABILIZATION POST-OPERATIVE GUIDELINES

### References

1. Andrews JR, Carson WG, Mcleod WD. Glenoid labrum tears related to the long head of the biceps. *Am J Sports Med.* 1985;13(5):337-341.
2. Fedoriw WW, Ramkumar P, Mcculloch PC, et al. Return to play after treatment of superior labral tears in. *Am J Sports Med.* 2014;42(5):1156-1159. doi:10.1177/0363546514528096.
3. Gehrman RM, Deluca PF, Bartolozzi AR. Humeral avulsion of the glenohumeral ligament caused by microtrauma to the anterior capsule in an overhand throwing athlete a case report. *Am J Sports Med.* 2003;31(4):617-619.
4. Gulotta L V, Lobatto D, Delos D, et al. Anterior shoulder capsular tears in professional baseball players. *J Shoulder Elb Surg.* 2018;23(8):e173-e178. doi:10.1016/j.jse.2013.11.027.
5. Kibler W Ben, Uhl TL, Maddux JWQ, et al. Qualitative clinical evaluation of scapular dysfunction: A reliability study. *J Shoulder Elb Surg.* 2002;11(6):550-556. doi:10.1067/mse.2002.126766.
6. Knesek M, Skendzel JG, Dines JS, et al. Diagnosis and management of superior labral anterior posterior tears in throwing athletes. *Am J Sports Med.* 2012;41(2):444-460. doi:10.1177/0363546512466067.
7. Taljanovic MS, Nisbet JK, Hunter TB, et al. Humeral avulsion of the inferior glenohumeral ligament in college female volleyball players caused by repetitive microtrauma. *Am J Sports Med.* 2011;39(5):1067-1076. doi:10.1177/0363546510391155.
8. Wilk KE, Arrigo CA, Hooks TR, et al. Rehabilitation of the overhead throwing athlete : there is more to it than just external rotation / internal rotation strengthening. *PMRJ.* 2016;8:78-90.
9. Wilk KE, MacRina LC, Arrigo C. Passive range of motion characteristics in the overhead baseball pitcher and their implications for rehabilitation. *Clin Orthop Relat Res.* 2012;470(6):1586-1594. doi:10.1007/s11999-012-2265-z.
10. Wilk KE, Obma P, Simpson CD, et al. Shoulder injuries in the overhead athlete. *J Orthop Sports Phys Ther.* 2009;39(2):38-54. doi:10.2519/jospt.2009.2929.
11. Wilk K, Yenchak AJ, Andrews JR. The Advanced Throwers Ten Exercise Program : A new exercise series for enhanced dynamic shoulder control in the overhead throwing athlete. *Phys Sportsmed.* 2011;39(4):90-97. doi:10.3810/psm.2011.11.1943.

12. Yoneda M, Nakagawa S, Hayashida K. Arthroscopic removal of symptomatic bennett lesions in the shoulders of baseball players : Arthroscopic. 2002;30(5):1-3.

Created: 1/2019

Reviewed: 3/2021; 4/2023