



Widespread Implementation of Medial Patellofemoral Ligament Reconstruction for Recurrent Patellar Instability Maintains Functional Outcomes at Midterm to Long-Term Follow-up While Decreasing Complication Rates: A Systematic Review

Kristen L. Stupay, B.A., Eric Swart, M.D., and Beth E. Shubin Stein, M.D.

Purpose: Our primary purpose was to evaluate whether complications have increased or functional outcomes have changed as medial patellofemoral ligament (MPFL) reconstruction has been adopted by more surgeons at more institutions over recent years. Our secondary purpose was to further define the complication profile of MPFL reconstruction. **Methods:** A systematic review of the literature was performed on January 12, 2014, using the keywords “medial patellofemoral ligament reconstruction,” “patellar instability reconstruction,” “patellofemoral ligament reconstruction,” and “MPFL.” Articles meeting our inclusion criteria were reviewed. Outcome measures, functional failures, complications, graft choice, and surgical technique were recorded and analyzed. **Results:** Thirty-four articles met our exclusion and inclusion criteria and were reviewed. Nineteen articles were “new” additions to the literature, whereas 15 had previously been reported on in prior analyses (“old”). The 19 new articles reported a statistically significant decrease in functional failure rates, from 9.55% in older studies to 4.77% in more recent studies ($P < .001$). The major complication rate dropped from 2.01% to 0.46% in the newer studies ($P = .005$), and the minor complication rate decreased from 6.53% to 4.00% ($P = .06$). Postoperative Kujala scores did not show a statistically significant change between newer and older publications (89.0 [SD, 3.7] and 89.4 [SD, 4.9], respectively; $P = .55$). Comparing results by fixation type, as well as by graft choice, showed no statistically significant differences in terms of outcomes or complication profile. **Conclusions:** With nearly twice the number of medical centers performing reconstruction of the MPFL and outcomes reported on nearly double the number of patients in recent years, functional outcomes remain favorable as complication and failure profiles are improving. Furthermore, despite a wide array of fixation techniques, as well as multiple options for graft constructs, there are no statistically or clinically significant differences in functional outcomes over time. This finding highlights the efficacy and adoptability of MPFL reconstruction for the treatment of recurrent patellar instability. **Level of Evidence:** Level IV, systematic review of mixed-level studies.

See commentary on page 1381

Recurrent patellar instability is a disabling condition that primarily affects young patients and often leads to cartilage injury, limiting patients’ participation

From Tulane University School of Medicine (K.L.S.), New Orleans, Louisiana; Columbia University Medical Center (E.S.); and the Department of Orthopaedic Surgery & Sports Medicine, Hospital for Special Surgery (B.E.S.S.), New York, New York, U.S.A.

The authors report that they have no conflicts of interest in the authorship and publication of this article.

Received July 27, 2014; accepted December 30, 2014.

Address correspondence to Kristen L. Stupay, B.A., Tulane University School of Medicine, 1430 Tulane Ave, New Orleans, LA 70112, U.S.A. E-mail: kstupay@tulane.edu

© 2015 by the Arthroscopy Association of North America
0749-8063/14650/\$36.00

<http://dx.doi.org/10.1016/j.arthro.2014.12.029>

in sports and potentially compromising their ability to carry out daily activities.¹⁻⁴ Furthermore, it is a particularly challenging problem for both the patient and the orthopaedic surgeon because of its multifactorial etiology. The most common contributing factors to recurrent patellar instability include bony variants, such as trochlear dysplasia, patella alta,⁵ and increased tibial tubercle–trochlear groove distance, as well as alignment issues, such as excessive genu valgum or tibial torsion.⁶ In addition to these predisposing bony conditions, incompetent soft-tissue stabilizers are implicated in the loss of patellar stability and can place the patient at high risk of recurrent dislocation.⁷

The reported overall recurrence rates after primary patellar dislocation range from 17%⁸ to more than

40%.^{9,10} After a second episode of instability, the recurrence rate rises to 50% or greater in some studies. Most investigators agree that this is the group in which surgical intervention should be considered¹¹ to minimize the risk of cartilage injury with subsequent dislocations. Surgical treatments for patellar instability are numerous and include both those aimed at reconstruction of the soft-tissue stabilizers (predominantly the medial patellofemoral ligament [MPFL]) and those that seek to correct malalignment and dysplasia through bony modifications, such as trochleoplasty or transfer of the tibial tubercle. In the past decade, there has been increased interest in ligamentous reconstructions to address soft tissues as an adjunct or alternative to bony corrections.

The primary soft-tissue restraint to lateral translation of the patella is the MPFL.¹²⁻¹⁵ The MPFL, which arises in the sulcus between the medial epicondyle and the adductor tubercle, runs deep and slightly distal to the vastus medialis and attaches to the proximal half of the medial aspect of the patella. Because of its limited ability to stretch, the MPFL is disrupted in most primary patellar dislocations and constitutes the dominant pathoanatomy associated with lateral patellar instability.^{7,14,16} In 1992 Ellera Gomes¹⁷ was the first author to propose that MPFL reconstruction was preferred over surgical correction of other predisposing conditions for treatment of recurrent patellar instability. This procedure has since been reported many times in the literature with good results, and it continues to gain popularity both as an isolated treatment and in combination with bony modifications for stabilization of the chronically dislocating patella.

The goal of MPFL reconstruction is to reconstitute the medial restraint of the patella, thereby resisting lateral dislocation and restoring stability. Many techniques are used to accomplish this, and various graft options have been described, including semitendinosus,¹⁸⁻²⁹ patellar tendon,^{30,31} partial quadriceps tendon,^{30,32} adductor magnus,^{6,30,33} gracilis,^{9,15,34-37} and iliotibial band grafts³⁸; allografts^{12,30,39}; and synthetic grafts.^{17,40,41} Methods for patellar fixation vary as well and include tunnels with interference screws^{12,22,39}; single^{40,42,43} and double^{34,44} through-and-through tunnels with looped, knotted,¹⁷ or soft-tissue suture fixation^{6,20,21}; V-shaped converging tunnels^{35,45}; double diverging tunnels²⁴; anchor fixation²³; and soft-tissue sutures.^{18,38} Fixation on the femoral side is most commonly achieved with a single blind tunnel and interference screw, but other methods include soft-tissue slings (particularly in skeletally immature patients),^{19,22} bone staples,^{40,43} through-and-through tunnels,³⁰ soft-tissue sutures,^{6,20,38} EndoButton (Smith & Nephew Endoscopy, Andover, MA) fixation,⁴⁶ and sutures tied over a bony bridge.²⁵

Systematic reviews following earlier outcomes of MPFL reconstruction included evaluation of radiologic outcomes,⁴⁷ functional outcomes,⁴⁸ rehabilitation and

return-to-sports efficacy,⁴⁹ and associated complications or failures.⁵⁰ The most recent systematic review addressing complications and failures, published by Shah et al.⁵⁰ in 2012, found that MPFL reconstruction was associated with a rather significant complication rate of 26.1%. They reported that 26 of the 597 patients included in their review actually had to return to the operating room for complications associated with their reconstruction, and of the 629 knees, 23 (3.7%) were deemed clinical failures at final follow-up. Clearly, despite the reporting of consistently positive clinical outcomes for most patients, the failure rate and complication profile of MPFL reconstruction demand further definition and continued improvement efforts.

In the past 3 years, the literature has seen rapid expansion in the number of studies reporting midterm results after MPFL reconstruction, as well as the number of different surgeons reporting outcomes. The primary purpose of this review was to address the question of whether, with increased adoption of this procedure as a treatment for recurrent instability, clinical outcomes have changed or complications have increased. We secondarily sought to further define the complication profile by asking whether any specific techniques are more prone to complications or poor outcomes.

On the basis of our experience with MPFL reconstruction for the treatment of recurrent patellar instability, we hypothesized that this procedure will continue to show good clinical outcomes as it is more widely used and it is followed up for longer periods. We further hypothesized that as more surgeons gain experience with this procedure and discover the particular surgical technique that works optimally for them and their patients, complications will decrease with time as well.

Methods

A systematic review of the literature was performed using the Medline, Embase, Scopus, and Web of Science databases. The purpose was to identify publications describing results of isolated MPFL reconstruction. The search was performed on January 12, 2014, and included publications that appeared in the literature as far back as 1992. The inclusion criteria were as follows: (1) The study must describe a technique for MPFL reconstruction in patients with chronic patellar instability with or without minor secondary soft-tissue surgery (e.g., release of the lateral retinaculum or advancement of the vastus medialis). (2) The study must require a minimum follow-up of 12 months in all enrolled patients. (3) The study article must be available in the English language. We excluded studies reporting MPFL reconstructions alongside other ligament reconstruction (anterior cruciate ligament, medial patellotibial ligament, and so on) or bony procedures, such as tibial tubercle osteotomies or trochleoplasty, were excluded. We also excluded

abstracts, animal studies, cadaveric studies, anatomic studies, biomechanical studies, case reports, technical notes, editorials, reviews, letters or comments to journals, and previous systematic reviews.

The databases were searched using the key terms “medial patellofemoral ligament reconstruction,” “patellar instability reconstruction,” “patellofemoral ligament reconstruction,” and “MPFL.” In addition, the reference lists of previous English-language systematic reviews addressing MPFL reconstruction were searched for publications that matched our criteria.⁴⁷⁻⁵⁰ The initial electronic search yielded 554 articles (250 in Medline, 59 in Embase, 166 in Scopus, and 79 in Web of Science). Once all duplicates were removed, there were 434 articles in total available for review. The titles and abstracts were assessed against the selection criteria, resulting in 64 potentially relevant publications. Full-text articles were then examined against the inclusion and exclusion criteria. At that time, 30 of the remaining articles were excluded for the following reasons: reporting a surgical procedure that was performed on patients with prior or concomitant bony modification (trochleoplasty or osteotomy); reporting on patients who did not meet the minimum follow-up of 12 months; reporting results of MPFL repair rather than reconstruction; and addressing the correction of acute, rather than recurrent, patellar dislocation. There were therefore 34 publications eligible for inclusion in this review, detailed in Tables 1 and 2. The search process is summarized in Figure 1.

In analyzing these publications, we were first and foremost interested in comparing functional clinical outcomes, as well as postoperative complications, in “old” versus “new” studies. We established a general cutoff for old and new studies based on the most recently published systematic review on MPFL reconstruction; all studies that were published before the literature search by Shah et al.⁵⁰ in October 2010 or that had been included in previous reviews were considered old. Any studies that were published after October 2010 or were not already included in another systematic review were considered new.

Our secondary interest was to look at the functional clinical outcomes and postoperative complications as they related to fixation techniques. A wide variety of surgical techniques were described in these articles. To make meaningful comparisons between groups, we broadly categorized the different reported techniques into the following: (1) femoral fixation technique, (2) patellar fixation technique, and (3) graft choice. Femoral and patellar fixation techniques were further divided between those that used bone tunnels or interference screws and those that used primarily soft-tissue attachments. Graft choices were separated into semitendinosus, gracilis, adductor magnus, patellar tendon, and synthetic ligament substitutes.

Functional clinical outcomes in all categories were compared using the Kujala Anterior Knee Pain Scale because it is a widely used and reliable score specific to knee function after MPFL reconstruction.⁵⁸ Postoperative complications were recorded for each publication and divided into (1) functional failures, (2) major complications, and (3) minor complications. Table 3 outlines the complications included in each category.

We made statistical comparisons between groups by comparing weighted outcomes using the Student 2-tailed *t* test (when comparing continuous outcomes such as Kujala scores). For comparisons of failure or complication rates, a χ^2 test on 2 proportions was used (or a Fisher exact test if the incidence was particularly low).

Results

Of the 34 publications included in this review, 15 had already been cited in prior systematic reviews whereas the remaining 19 represented new contributions to the literature. Of the 19 new studies, 18 were published by medical centers that had not previously reported outcomes with this procedure (31 total centers included) and represented an additional 621 patients and 650 knees (of 1,015 patients and 1,048 knees in total).

A comparison of our results by date of publication is shown in Table 4. The mean Kujala scores were 89.4 (SD, 4.9) for old publications and 89.0 (SD, 3.7) for new publications, which was not statistically significant ($P = .55$). The overall functional failure rate was 9.55% (38 of 398) in older publications and dropped to 4.77% (31 of 650) in the newer series, which was statistically significant ($P < .001$). The major complication rate dropped from 2.01% (8 of 398) in older studies to 0.46% (3 of 650) in newer studies, which was statistically significant ($P = .005$). The minor complication rate also decreased between the older and newer studies (6.53% and 4.00%, respectively); however, this finding was not significant ($P = .06$). These findings are summarized in Table 5.

Comparing results by fixation type showed no statistically significant differences in terms of functional outcomes or complication profile. For patellar fixation, 21 studies used bone tunnels, with a mean Kujala score of 88.6 (SD, 4.7), whereas 10 groups used only soft-tissue attachments, with a mean Kujala score of 90.0 (SD, 3.7), which was not significantly different ($P = .38$). For femoral fixation, 24 studies reported outcomes using bone tunnels whereas 7 used soft-tissue techniques to attach the graft, which was also not significantly different (Kujala scores of 88.5 [SD, 4.0] and 93.8 [SD, 3.2], respectively; $P = .06$). Regarding graft choice, 17 studies used semitendinosus grafts, 3 used gracilis grafts, and 3 used artificial grafts whereas the remaining 11 had heterogeneous graft choices. Kujala scores were similar between semitendinosus and

Table 1. New Articles, Published After October 2010, or Not Previously Reviewed

| Authors | Year | No. of Knees | Mean Age, yr | Mean Follow-up, mo | Patellar Fixation | Femoral Fixation | Graft | Functional Failures, n | Complications, n | | Mean Kujala Score | |
|-----------------------------------|------|--------------|--------------|--------------------|-------------------|------------------|-------|------------------------|------------------|-------|-------------------|---------------|
| | | | | | | | | | Major | Minor | Preoperative | Postoperative |
| Csintalan et al. ⁵¹ | 2014 | 56 | 24 | 51.6 | BT | BT | ST | 6 | 0 | 5 | — | — |
| Deie et al. ¹⁸ | 2011 | 31 | 22.2 | 38.4 | STA | BT | ST | 1 | 0 | 1 | 64 | 94.5 |
| Goyal et al. ⁵² | 2013 | 32 | 25 | 38 | STA | BT | QU | 0 | 0 | 0 | 49.31 | 91.25 |
| Hinterwimmer et al. ⁵³ | 2013 | 19 | 23 | 16 | BT | BT | GR | 0 | 1 | 2 | — | 92.7 |
| Kang et al. ²¹ | 2013 | 82 | 28.86 | 14 | STA | BT | ST | 0 | 0 | 0 | 52.98 | 93.54 |
| Kumahashi et al. ²² | 2012 | 5 | 13.6 | 27.8 | BT | STA | ST | 0 | 0 | 0 | 67.4 | 95.4 |
| Ma et al. ²³ | 2013 | 32 | 28.4 | 40 | STA | BT | ST | 0 | 0 | 0 | 54 | 87 |
| Nelitz et al. ³⁵ | 2013 | 21 | 12.2 | 33.6 | BT | BT | GR | 2 | 0 | 0 | 72.9 | 92.8 |
| Panni et al. ²⁴ | 2011 | 48 | 28 | 33 | BT | BT | ST | 0 | 1 | 6 | 56.7 | 86.8 |
| Raghuveer et al. ⁵⁴ | 2012 | 15 | 29.2 | 42 | Mixed | BT | Mixed | 2 | 0 | 5 | 44.8 | 91.9 |
| Slenker et al. ³⁹ | 2013 | 35 | 20.6 | 21 | BT | BT | Mixed | 3 | 0 | 3 | 49 | 89.5 |
| Sobhy et al. ²⁵ | 2013 | 29 | 20.1 | 32.2 | BT | BT | ST | 0 | 1 | 0 | 36.6 | 90.6 |
| Song et al. ²⁶ | 2014 | 20 | 21 | 34.5 | SA | BT | ST | 1 | 0 | 0 | 52.6 | — |
| Wagner et al. ⁵⁵ | 2013 | 50 | 19 | 12 | STA | BT | GR | 1 | 0 | 2 | 70 | 87 |
| Wang et al. ³⁷ | 2013 | 70 | 25 | 48 | SA | BT | ST | 3 | 0 | 2 | 48.9 | 88.25 |
| Wang et al. ²⁷ | 2010 | 69 | 30.2 | 42 | BT | BT | ST | 11 | 0 | 0 | 52.73 | 82.27 |
| Wang et al. ²⁸ | 2012 | 22 | 23 | 37.5 | STA | BT | ST | 1 | 0 | 0 | 53.9 | 84.1 |
| Witonski et al. ⁴⁵ | 2013 | 10 | 27.2 | 43 | STA* | SA | PT | 0 | 0 | 0 | 59.7 | 84.4 |
| Yercan et al. ²⁹ | 2011 | 4 | 8.7 | 17.7 | BT | STA | ST | 0 | 0 | 0 | 36 | 89.5 |

BT, bone tunnel; GR, gracilis; PT, patellar tendon; QU, quadriceps; SA, suture anchor; ST, semitendinosus; STA, soft-tissue attachment. *Patellar tendon remained at natural insertion.

gracilis grafts (89.0 [SD, 4.9] and 89.6 [SD, 3.3], respectively; *P* = .8). Only one study using artificial grafts reported Kujala scores, so statistical comparisons were not possible. These results are summarized in Table 6.

Discussion

The purpose of this review was to evaluate the midterm to long-term functional outcomes of the various surgical techniques that are in use for reconstruction of the MPFL. Surely, knowledge gained from early studies, refined techniques, and improvements in

the rehabilitation process over recent years have affected the outcomes for the better; however, with the adoption of these new techniques across increasingly large physician and patient populations, it is necessary to consider whether the results of the initial pioneers are reproducible in the general orthopaedic community. Therefore we also sought to assess whether the rapid adoption of MPFL reconstruction has altered the success or complication rates that were seen in the earlier literature.

For this review, we identified 34 articles, published between 1992 and January 2014, reporting clinical

Table 2. Old Articles, Published Before October 2010, or Previously Reviewed

| Authors | Year | No. of Knees | Mean Age, yr | Mean Follow-up, mo | Patellar Fixation | Femoral Fixation | Graft | Functional Failures, n | Complications, n | | Mean Kujala Score | |
|-----------------------------------|------|--------------|--------------|--------------------|-------------------|------------------|-------|------------------------|------------------|-------|-------------------|---------------|
| | | | | | | | | | Major | Minor | Preoperative | Postoperative |
| Ahmad et al. ¹² | 2009 | 20 | 23 | 31 | BT | BT | Mixed | 0 | 0 | 1 | 49.9 | 88.2 |
| Deie et al. ¹⁹ | 2005 | 46 | 19.2 | 114 | BT | STA | ST | 8 | 0 | 0 | 50 | 95 |
| Drez et al. ³⁸ | 2001 | 14 | 22 | 31.5 | STA | STA | Mixed | 1 | 1 | 2 | — | 88.6 |
| Ellera Gomes ¹⁷ | 1992 | 30 | 29 | 39 | BT | BT | PLY | 1 | 1 | 3 | — | — |
| Ellera Gomes et al. ²⁰ | 2004 | 16 | 26.7 | 60 | BT | BT | ST | 1 | 1 | 1 | — | — |
| Gomes ⁶ | 2008 | 24 | 19.3 | 53 | BT | BT | Mixed | 1 | 0 | 0 | — | — |
| Han et al. ⁴⁴ | 2011 | 59 | 24.3 | 68.4 | BT | BT | ST | 0 | 0 | 3 | 41.4 | 82.6 |
| Nomura et al. ⁴⁰ | 2000 | 27 | 21 | 70.8 | BT | STA | PLY | 4 | 0 | 4 | — | — |
| Nomura and Inoue ⁴² | 2006 | 12 | 24.8 | 50.4 | BT | STA | ST | 0 | 0 | 3 | 56.3 | 96 |
| Nomura et al. ⁴³ | 2007 | 24 | 22.5 | 142.8 | BT | STA | PLY | 7 | 0 | 1 | 63.2 | 94.2 |
| Ronga et al. ⁵⁶ | 2009 | 28 | 32.5 | 37.2 | BT | BT | Mixed | 3 | 0 | 4 | 45 | 83 |
| Sillanpää et al. ³³ | 2008 | 15 | 20.2 | 121.2 | STA | — | AM | 3 | 1 | 0 | — | 88 |
| Steiner et al. ³⁰ | 2006 | 34 | 27 | 66.5 | BT | BT | Mixed | 0 | 1 | 4 | 53.3 | 90.7 |
| Toritsuka et al. ⁴⁶ | 2011 | 20 | 23 | 30 | BT | BT | ST | 1 | 1 | 0 | — | 96 |
| Watanabe et al. ⁵⁷ | 2008 | 29 | 19 | 51.6 | STA | EB | Mixed | 8 | 2 | 0 | — | — |

AM, adductor magnus; BT, bone tunnel; EB, EndoButton; PLY, polyester; ST, semitendinosus; STA, soft-tissue attachment.



Fig 1. Search process. (f/u, follow-up; MPFL, medial patellofemoral ligament; SR, systematic review; Tech, technical; WOS, Web of Science.)

outcomes of MPFL reconstruction for recurrent patellar instability, 19 of which represent new contributions to the literature since October 2010. These 19 newer articles were published by 18 medical centers that had not previously produced literature on MPFL reconstruction and included 621 patients, which represents more than double the total number of patients on whom results were reported in the articles published before October 2010. Our analysis of the data from these 34 publications indicates that MPFL reconstruction has maintained very good functional outcomes since first being published in 1992. The mean postoperative Kujala score for all 1,015 patients included in this review was 89.1, up from a mean preoperative score of 53.1; this was observed alongside a decrease in the failure rate from 9.55% in studies published during or before October 2010 to 4.77% in newer

studies. Furthermore, the major postoperative complication rate in the newer studies was significantly lower than that in the older studies. Lastly, we found that soft-tissue fixation techniques did not statistically differ from bone tunnel fixation techniques regarding functional outcomes, nor did the graft choice seem to alter postoperative function of the reconstructed ligament. Thus what we have seen through critical appraisal of the data is encouraging: More physicians in more medical centers are performing this procedure on more patients than before, are doing so with equal functional success, and yet seem to be encountering fewer postoperative failures and complications regardless of fixation technique or graft used.

Existing literature on MPFL reconstruction has come to complementary conclusions. In 2010 Buckens and

Table 3. Postoperative Failures and Complications

| Functional Failures | Major Complications | Minor Complications |
|-------------------------------------|------------------------------------|-------------------------------------|
| Clinical apprehension sign | Patellar fracture | ROM deficit >10°, corrected |
| Patient-reported repeat subluxation | ROM deficit >10°, uncorrected | Stiffness requiring MUA |
| Patient-reported repeat dislocation | RTOR because of graft complication | Persistent pain |
| | Unable to run | RTOR for HR |
| | | Superficial wound infection |
| | | Extensor lag |
| | | Wound complications, noninfectious* |
| | | Subcutaneous hematoma |

HR, hardware removal; MUA, manipulation under anesthesia; ROM, range of motion; RTOR, return to operating room.

*Dehiscence and trouble healing requiring a return to the operating room.

Table 4. Postoperative Kujala Scores, Functional Failure Rates, and Complication Rates in Old Versus New Studies

| | No. of Studies | No. of Knees | Mean Postoperative Kujala Score | Functional Failures, % | Major Complications, % | Minor Complications, % |
|----------------|----------------|--------------|---------------------------------|------------------------|------------------------|------------------------|
| New studies | 19 | 650 | 89.0 (SD, 3.7) | 4.77% | 0.46% | 4.00% |
| Old studies | 15 | 398 | 89.4 (SD, 4.9) | 9.55% | 2.01% | 6.53% |
| <i>P</i> value | | | .55 | < .001 | .005 | .06 |

Saris⁴⁸ evaluated MPFL reconstruction alongside other realignment and stabilization techniques, such as osteotomy or trochleoplasty. They critically reviewed 21 publications and, although their appraisal did not produce any statistically significant outcomes, were able to show a trend that MPFL reconstruction provided equal, if not superior, functional outcomes to alleviate pain and realign the patella, with lower perioperative morbidity rates and fewer long-term complications as compared with older techniques. Fisher et al.⁴⁹ in 2010 asserted that MPFL reconstruction is very likely to improve a patient’s ability to perform routine activities of daily living, regardless of the particular reconstruction method or graft construct used to perform the procedure. This finding is consistent with ours and supports the notion that the surgeon may choose whichever graft is best suited for the individual patient, without concern for compromising outcomes. Shah et al.⁵⁰ in 2012 took on the task of trying to delineate complication rates associated with the myriad techniques currently in use for reconstruction of the MPFL—namely, tunnel fixation versus soft-tissue and/or suture anchor fixation. They did not identify any clear statistical relation among techniques, suggesting that the different techniques used to fix the reconstructed MPFL are all equally reliable as long as sound mechanical principles are observed. It would therefore

seem that when reconstructing the MPFL, the surgeon may choose whichever method of fixation with which he or she is most comfortable. In 2013 Vavken et al.⁵⁹ concluded that, in the pediatric and adolescent population, MPFL reconstruction is the most effective treatment option for chronic patellar instability and is safe regarding physis growth and development in the young patient. These findings indicate that MPFL reconstruction can be safely and reliably implemented in any age group, which further supports the use of MPFL reconstruction in an ever-broadening population of patients.

Limitations

In writing this review, we did encounter some challenges that limit the generalizability of the results. There is a paucity of high-level evidence in the current body of literature on patellar instability; only 2 studies were classified as Level II studies, and we identified no Level I studies. Methodologic deficiencies regarding sample size, follow-up period, and varying adjunctive surgical procedures obscured our ability to draw absolute conclusions from the data collected. We attempted to minimize confounding effects of additional stabilizing procedures by excluding the studies reporting patients who had undergone adjunctive procedures that required bony modifications—such as osteotomy and trochleoplasty—or who had undergone additional

Table 5. Complication Profile in Old Versus New Studies

| | No. of Studies | No. of Knees | Major Complications | Minor Complications |
|----------------|----------------|--------------|--|--|
| New studies | 19 | 650 | 3 of 650, 0.46% 2 with patellar fracture 1 with ROM deficit >10°, uncorrected | 26 of 650, 4.00% 9 with persistent knee pain 4 with ROM deficit >10°, corrected 4 with RTOR for hardware removal 4 with superficial wound infections 2 with extensor lag 2 with wound complications, noninfectious* 1 with stiffness requiring MUA |
| Old studies | 15 | 398 | 8 of 398, 2.01% 3 with RTOR for graft complications (fibrous bridge, dislocation, loosening) 1 with patellar fracture 1 unable to run postoperatively | 26 of 398, 6.53% 6 with persistent knee pain 4 with stiffness requiring IPT 4 with wound complications, noninfectious 3 with ROM deficit >10°, corrected 3 with stiffness requiring MUA 3 with RTOR for hardware removal 2 with subcutaneous hematoma |
| <i>P</i> value | | | <i>P</i> = .005 | <i>P</i> = .06 |

IPT, intensive physical therapy; MUA, manipulation under anesthesia; ROM, range of motion; RTOR, return to operating room.
*Dehiscence and trouble healing requiring a return to the operating room.

Table 6. Postoperative Kujala Scores Based on Patellar and Femoral Fixation Technique and Graft Type

| | Postoperative Kujala Score |
|----------------------------|----------------------------|
| Bone tunnel | |
| Patellar fixation (n = 21) | 88.6 (4.7) |
| Femoral fixation (n = 24) | 88.5 (4.0) |
| Soft tissue | |
| Patellar fixation (n = 10) | 90.0 (3.7) |
| Femoral fixation (n = 7) | 93.8 (3.2) |
| Graft type | |
| Semitendinosus (n = 17) | 89.0 (4.9) |
| Gracilis (n = 3) | 89.6 (3.3) |

NOTE. Data are presented as mean (standard deviation).

ligament reconstructions. We did not, however, exclude studies whose patients had undergone soft-tissue procedures, such as lateral retinaculum release, chondroplasty, or advancement of the vastus medialis obliquus. Of the 34 studies that we included, 25 included patients who underwent at least 1 of the aforementioned soft-tissue procedures either before or concomitantly with MPFL reconstruction. In addition, we hoped to improve validity by requiring a minimum patient follow-up of 12 months, a requirement that has not been made in prior systematic reviews to this point. Regardless, it is possible that postoperative complications and/or failures were missed as a result of not demanding a longer follow-up period.

We encountered another issue in analyzing the major and minor complication rates as a function of the type of graft fixation used because reporting standards are variable throughout the literature. Because most of these studies are retrospective, they are also associated with the usual issues revolving around recall bias and the like. This problem has been present in previous reviews of MPFL reconstruction literature, and we believe that the literature could benefit from more uniform and transparent reporting of complications, ideally in a prospective manner. Lastly, we excluded 17 articles whose full texts were not available in the English language. Despite these limitations, we believe that meaningful conclusions may still be gleaned from the existing literature.

Conclusions

With nearly twice the number of medical centers performing reconstruction of the MPFL and outcomes reported on nearly double the number of patients in recent years, functional outcomes remain favorable as complication and failure profiles are improving. Furthermore, despite a wide array of fixation techniques, as well as multiple options for graft constructs, there are no statistically or clinically significant differences in functional outcomes over time. This finding highlights the efficacy and adoptability of MPFL reconstruction for the surgical treatment of recurrent

patellar instability and bodes well for generalized adoption of this procedure by the orthopaedic community. Still, the treatment of recurrent patellar instability warrants further investigation in the future, with larger, prospective, high-quality trials that include more reliable and uniform reporting of postoperative complications.

References

- Mäenpää H, Lehto MU. Patellar dislocation. The long-term results of nonoperative management in 100 patients. *Am J Sports Med* 1997;25:213-217.
- Camanho GL, Viegas Ade C, Bitar AC, Demange MK, Hernandez AJ. Conservative versus surgical treatment for repair of the medial patellofemoral ligament in acute dislocations of the patella. *Arthroscopy* 2009;25:620-625.
- Mäenpää H, Huhtala H, Lehto MU. Recurrence after patellar dislocation. Redislocation in 37/75 patients followed for 6-24 years. *Acta Orthop Scand* 1997;68:424-426.
- Petri M, Liதாகி E, Hofmeister M, et al. Operative vs conservative treatment of traumatic patellar dislocation: Results of a prospective randomized controlled clinical trial. *Arch Orthop Trauma Surg* 2013;133:209-213.
- Dejour D, Le Coultre B. Osteotomies in patello-femoral instabilities. *Sports Med Arthrosc Rev* 2007;15:39-46.
- Gomes JE. Comparison between a static and a dynamic technique for medial patellofemoral ligament reconstruction. *Arthroscopy* 2008;24:430-435.
- Nomura E, Horiuchi Y, Inoue M. Correlation of MR imaging findings and open exploration of medial patellofemoral ligament injuries in acute patellar dislocations. *Knee* 2002;9:139-143.
- Fithian DC, Paxton EW, Stone ML, et al. Epidemiology and natural history of acute patellar dislocation. *Am J Sports Med* 2004;32:1114-1121.
- Marteau E, Burdin P, Brillhault J. Gracilis tendon transfer associated with distal alignment for patella alta with recurrent dislocations: An original surgical technique. *Orthop Traumatol Surg Res* 2011;97:S5-S11 (suppl).
- Smith TO, Song F, Donell ST, Hing CB. Operative versus non-operative management of patellar dislocation. A meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 2011;19:988-998.
- Lind M, Jakobsen BW, Lund B, Christiansen SE. Reconstruction of the medial patellofemoral ligament for treatment of patellar instability. *Acta Orthop* 2008;79:354-360.
- Ahmad CS, Brown GD, Stein BS. The docking technique for medial patellofemoral ligament reconstruction: Surgical technique and clinical outcome. *Am J Sports Med* 2009;37:2021-2027.
- Panagopoulos A, van Niekerk L, Triantafillopoulos IK. MPFL reconstruction for recurrent patella dislocation: A new surgical technique and review of the literature. *Int J Sports Med* 2008;29:359-365.
- Desio SM, Burks RT, Bachus KN. Soft tissue restraints to lateral patellar translation in the human knee. *Am J Sports Med* 1998;26:59-65.
- Conlan T, Garth WP Jr, Lemons JE. Evaluation of the medial soft-tissue restraints of the extensor mechanism of the knee. *J Bone Joint Surg Am* 1993;75:682-693.

16. Sallay PI, Poggi J, Speer KP, Garrett WE. Acute dislocation of the patella. A correlative pathoanatomic study. *Am J Sports Med* 1996;24:52-60.
17. Ellera Gomes JL. Medial patellofemoral ligament reconstruction for recurrent dislocation of the patella: A preliminary report. *Arthroscopy* 1992;8:335-340.
18. Deie M, Ochi M, Adachi N, Shibuya H, Nakamae A. Medial patellofemoral ligament reconstruction fixed with a cylindrical bone plug and a grafted semitendinosus tendon at the original femoral site for recurrent patellar dislocation. *Am J Sports Med* 2011;39:140-145.
19. Deie M, Ochi M, Sumen Y, Adachi N, Kobayashi K, Yasumoto M. A long-term follow-up study after medial patellofemoral ligament reconstruction using the transferred semitendinosus tendon for patellar dislocation. *Knee Surg Sports Traumatol Arthrosc* 2005;13:522-528.
20. Ellera Gomes JL, Stigler Marczyk LR, Cesar de Cesar P, Jungblut CF. Medial patellofemoral ligament reconstruction with semitendinosus autograft for chronic patellar instability: A follow-up study. *Arthroscopy* 2004;20:147-151.
21. Kang H, Cao J, Yu D, Zheng Z, Wang F. Comparison of 2 different techniques for anatomic reconstruction of the medial patellofemoral ligament: A prospective randomized study. *Am J Sports Med* 2013;41:1013-1021.
22. Kumahashi N, Kuwata S, Tadenuma T, Kadowaki M, Uchio Y. A "sandwich" method of reconstruction of the medial patellofemoral ligament using a titanium interference screw for patellar instability in skeletally immature patients. *Arch Orthop Trauma Surg* 2012;132:1077-1083.
23. Ma LF, Wang F, Chen BC, Wang CH, Zhou JW, Wang HY. Medial retinaculum plasty versus medial patellofemoral ligament reconstruction for recurrent patellar instability in adults: A randomized controlled trial. *Arthroscopy* 2013;29:891-897.
24. Panni AS, Alam M, Cerciello S, Vasso M, Maffulli N. Medial patellofemoral ligament reconstruction with a divergent patellar transverse 2-tunnel technique. *Am J Sports Med* 2011;39:2647-2655.
25. Sobhy MH, Mahran MA, Kamel EM. Midterm results of combined patellofemoral and patellotibial ligaments reconstruction in recurrent patellar dislocation. *Eur J Orthop Surg Traumatol* 2013;23:465-470.
26. Song SY, Kim IS, Chang HG, Shin JH, Kim HJ, Seo YJ. Anatomic medial patellofemoral ligament reconstruction using patellar suture anchor fixation for recurrent patellar instability. *Knee Surg Sports Traumatol Arthrosc* 2014;22:2431-2437.
27. Wang F, Kang HJ, Chen BC, Chen W, Su YL, Zhang YZ. Combination of medial patellofemoral ligament reconstruction with vastus medialis advancement for chronic patellar dislocation. *Chin Med J (Engl)* 2010;123:3024-3029.
28. Wang JL, Li HP, Liu YJ, et al. Reconstruction of the medial patellofemoral ligament with a suture-tie technique of patellar side fixation. *Chin Med J (Engl)* 2012;125:1884-1888.
29. Yercan HS, Erkan S, Okcu G, Ozalp RT. A novel technique for reconstruction of the medial patellofemoral ligament in skeletally immature patients. *Arch Orthop Trauma Surg* 2011;131:1059-1065.
30. Steiner TM, Torga-Spak R, Teitge RA. Medial patellofemoral ligament reconstruction in patients with lateral patellar instability and trochlear dysplasia. *Am J Sports Med* 2006;34:1254-1261.
31. Camanho GL, Bitar AC, Hernandez AJ, Olivi R. Medial patellofemoral ligament reconstruction: A novel technique using the patellar ligament. *Arthroscopy* 2007;23:108.e1-108.e4.
32. Dopirak R, Adamany D, Bickel B, Steensen R. Reconstruction of the medial patellofemoral ligament using a quadriceps tendon graft: A case series. *Orthopedics* 2008;31:217.
33. Sillanpää P, Mattila VM, Visuri T, Mäenpää H, Pihlajamäki H. Ligament reconstruction versus distal realignment for patellar dislocation. *Clin Orthop Relat Res* 2008;466:1475-1484.
34. Hinterwimmer S, Imhoff AB, Minzlaff P, et al. Anatomical two-bundle medial patellofemoral ligament reconstruction with hardware-free patellar graft fixation: Technical note and preliminary results. *Knee Surg Sports Traumatol Arthrosc* 2013;21:2147-2154.
35. Nelitz M, Dreyhaupt J, Reichel H, Woelfle J, Lippacher S. Anatomic reconstruction of the medial patellofemoral ligament in children and adolescents with open growth plates: Surgical technique and clinical outcome. *Am J Sports Med* 2013;41:58-63.
36. Siebold R, Borbon CAV. Arthroscopic extraarticular reconstruction of the medial patellofemoral ligament with gracilis tendon autograft—Surgical technique. *Knee Surg Sports Traumatol Arthrosc* 2012;20:1245-1251.
37. Wang CH, Ma LF, Zhou JW, et al. Double-bundle anatomical versus single-bundle isometric medial patellofemoral ligament reconstruction for patellar dislocation. *Int Orthop* 2013;37:617-624.
38. Drez D Jr, Edwards TB, Williams CS. Results of medial patellofemoral ligament reconstruction in the treatment of patellar dislocation. *Arthroscopy* 2001;17:298-306.
39. Slenker NR, Tucker BS, Pepe MD, Marchetto PA, Cohen SB. Short-/intermediate-term outcomes after medial patellofemoral ligament reconstruction in the treatment of chronic lateral patellofemoral instability. *Phys Sportsmed* 2013;41:26-33.
40. Nomura E, Horiuchi Y, Kihara M. A mid-term follow-up of medial patellofemoral ligament reconstruction using an artificial ligament for recurrent patellar dislocation. *Knee* 2000;7:211-215.
41. Nomura E, Inoue M. Surgical technique and rationale for medial patellofemoral ligament reconstruction for recurrent patellar dislocation. *Arthroscopy* 2003;19:E47.
42. Nomura E, Inoue M. Hybrid medial patellofemoral ligament reconstruction using the semitendinous tendon for recurrent patellar dislocation: Minimum 3 years' follow-up. *Arthroscopy* 2006;22:787-793.
43. Nomura E, Inoue M, Kobayashi S. Long-term follow-up and knee osteoarthritis change after medial patellofemoral ligament reconstruction for recurrent patellar dislocation. *Am J Sports Med* 2007;35:1851-1858.
44. Han H, Xia Y, Yun X, Wu M. Anatomical transverse patella double tunnel reconstruction of medial patellofemoral ligament with a hamstring tendon autograft for

- recurrent patellar dislocation. *Arch Orthop Trauma Surg* 2011;131:343-351.
45. Witonski D, Keska R, Synder M, Sibinski M. An isolated medial patellofemoral ligament reconstruction with patellar tendon autograft. *Biomed Res Int* 2013;2013:637678.
 46. Toritsuka Y, Amano H, Mae T, et al. Dual tunnel medial patellofemoral ligament reconstruction for patients with patellar dislocation using a semitendinosus tendon autograft. *Knee* 2011;18:214-219.
 47. Smith TO, Russell N, Walker J. A systematic review investigating the early rehabilitation of patients following medial patellofemoral ligament reconstruction for patellar instability. *Crit Rev Phys Rehabil Med* 2007;19:79-95.
 48. Buckens CFM, Saris DBF. Reconstruction of the medial patellofemoral ligament for treatment of patellofemoral instability: A systematic review. *Am J Sports Med* 2010;38:181-188.
 49. Fisher B, Nyland J, Brand E, Curtin B. Medial patellofemoral ligament reconstruction for recurrent patellar dislocation: A systematic review including rehabilitation and return-to-sports efficacy. *Arthroscopy* 2010;26:1384-1394.
 50. Shah JN, Howard JS, Flanigan DC, Brophy RH, Carey JL, Lattermann C. A systematic review of complications and failures associated with medial patellofemoral ligament reconstruction for recurrent patellar dislocation. *Am J Sports Med* 2012;40:1916-1923.
 51. Csintalan RP, Latt LD, Fornalski S, Raiszadeh K, Inacio MC, Fithian DC. Medial patellofemoral ligament (MPFL) reconstruction for the treatment of patellofemoral instability. *J Knee Surg* 2014;27:139-146.
 52. Goyal D. Medial patellofemoral ligament reconstruction: The superficial quad technique. *Am J Sports Med* 2013;41:1022-1029.
 53. Hinterwimmer S, Imhoff AB, Minzlaff P, et al. Anatomical two-bundle medial patellofemoral ligament reconstruction with hardware free patellar graft fixation: A technical note and preliminary results. *Knee Surg Sports Traumatol Arthrosc* 2013;21:2147-2154.
 54. Raghuvveer RK, Mishra CB. Reconstruction of medial patellofemoral ligament for chronic patellar instability. *Indian J Orthop* 2012;46:447-454.
 55. Wagner D, Pfalzer F, Hingelbaum S, Huth J, Mauch F, Bauer G. The influence of risk factors on clinical outcomes following anatomical medial patellofemoral ligament (MPFL) reconstruction using the gracilis tendon. *Knee Surg Sports Traumatol Arthrosc* 2013;21:318-324.
 56. Ronga M, Oliva F, Longo UG, Testa V, Capasso G, Maffulli N. Isolated medial patellofemoral ligament reconstruction for recurrent patellar dislocation. *Am J Sports Med* 2009;37:1735-1742.
 57. Watanabe T, Muneta T, Ikeda H, Tateishi T, Sekiya I. Visual analog scale assessment after medial patellofemoral ligament reconstruction with or without tibial tubercle transfer. *J Orthop Sci* 2008;13:32-38.
 58. Kujala UM, Jaakkola LH, Koskinen SK, Taimela S, Hurme M, Nelimarkka O. Scoring of patellofemoral disorders. *Arthroscopy* 1993;9:159-163.
 59. Vavken P, Wimmer MD, Camathias C, Quidde J, Valderrabano V, Pagenstert G. Treating patella instability in skeletally immature patients. *Arthroscopy* 2013;29:1410-1422.