

SPECIALTY UPDATE

What's New in Limb Lengthening and Deformity Correction

Reggie C. Hamdy, MD, MSc, FRCS(C), Mitchell Bernstein, MD, FRCSC, Austin T. Fragomen, MD, and S. Robert Rozbruch, MD

Investigation performed at the Shriners Hospital for Children, Montreal, Quebec, Canada

Limb reconstruction surgery continues to affirm itself as a distinct subspecialty worldwide. Apart from the specialty day at the annual American Academy of Orthopaedic Surgeons (AAOS) meeting, the annual Limb Lengthening and Reconstruction Society (LLRS) North American meeting, the combined Association for the Study and Application of the Methods of Ilizarov (ASAMI)/International LLRS (ASAMI/ILLRS) meeting in Brisbane, Queensland, Australia, and the very popular annual deformity courses (Baltimore, Hull, Cairo, and others), national ASAMI meetings were held in many countries. Two new issues of the *Journal of Limb Lengthening & Reconstruction* were published in 2016 with 20 manuscripts, including editorials on why deformity concepts are still not a mainstream part of orthopaedics¹, the evolution of limb lengthening and reconstruction in the post-Ilizarov era², and what happened to the regenerate following the numerous modifications of the original Ilizarov technique³.

Pediatrics

Congenital Limb Deficiencies

A most comprehensive description of surgical treatment of tibial and fibular hemimelia was reported by Paley^{4,5}. Eidelman et al. recommended a routine performance of pelvic osteotomy for patients with Paley Type 1B deficiency undergoing lengthening⁶. Saldanha et al. stressed the importance of systematic assessment of the knee prior to any intervention in children with congenital limb deficiencies⁷. Foot preservation and reconstruction of tibial hemimelia were reported to provide good functional outcome⁸. The Ilizarov method provided comparable results in the treatment of congenital pseudarthrosis of the tibia whether it was associated with neurofibromatosis type 1 (NF1)

or was idiopathic in origin in pediatric cases⁹. The use of the Fassier-Duval rod as an adjunct procedure in the treatment of congenital pseudarthrosis of the tibia was shown to be successful in obtaining union in 4 children¹⁰. A new successful technique for the treatment of congenital pseudarthrosis of the tibia was described¹¹.

Pediatric Trauma

The role of hexapod external fixation in the treatment of tibial fractures in children was discussed by Iobst¹². Ilizarov bone transport and fibular bone graft for reconstruction of segmental tibial defects were reported to give equally effective results¹³. Shore et al. compared the cost and complications of uniplanar external fixation with those of Taylor Spatial Frame (Smith & Nephew) external fixation in the treatment of pediatric diaphyseal fractures and found that a corrected cost analysis revealed equivalent costs for care delivery¹⁴.

Limb-Length Discrepancy

Mills and Nelson reported on an improved spreadsheet for calculating limb-length discrepancy and epiphysiodesis timing using the multiplier method¹⁵. Poor efficiency and unpredictability of epiphysiodesis using Eight-Plates (Orthofix) for the treatment of limb-length discrepancy were demonstrated by Gaumétou et al.¹⁶. Makarov et al. reported a 7% complication rate, with angular deformity being the most common complication, in 863 children who underwent epiphysiodesis for the treatment of limb-length discrepancy¹⁷. Donnan et al. reported that a 2-ring tibial lengthening fixator was effective in maintaining segmental alignment and callus production during tibial lengthening¹⁸. In a comparative study evaluating 2 techniques of syndesmotic fixation, transverse tricortical screws and oblique quadricortical screws, the authors found no advantage of 1 technique over the other¹⁹. Yoshida et al. evaluated the

Specialty Update has been developed in collaboration with the Board of Specialty Societies (BOS) of the American Academy of Orthopaedic Surgeons.

Disclosure: One author of this work (R.C.H.) received a stipend from JBJS for writing this work. On the **Disclosure of Potential Conflicts of Interest** forms, which are provided with the online version of the article, one or more of the authors checked "yes" to indicate that the author had a relevant financial relationship in the biomedical arena outside the submitted work (<http://links.lww.com/JBJS/E312>).

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

complications related to fibular resection during tibial lengthening performed with a Taylor Spatial Frame and recommended fixing the tibial fibular joints by transfixing the wire and/or cannulated screws both proximally and distally to minimize proximal and distal fibular migration²⁰.

Skeletal Dysplasias

Extensive limb lengthening of >50% of the initial length in patients with achondroplasia or hypochondroplasia was reported to carry a considerable risk of complications²¹. Knee malalignment was evaluated in 581 children with Down syndrome by Duque Orozco et al., who found that 5% had patellofemoral instability and other underlying knee deformities²², a relevant finding in situations in which limb lengthening is considered. Distraction osteogenesis appears to be an effective and safe approach for the simultaneous correction of the shortness of the first ray as well as the medial angulation of the great toe in patients with Apert syndrome²³. The mean tibial slope in patients with achondroplasia was found to be substantially more anterior, and this may predispose patients to genu recurvatum²⁴.

Growth Modulation

Stevens presented an excellent review on various strategies using reversible and serial guided growth in limb lengthening²⁵. Farr et al. reported a high rate of radiographic recurrence of frontal plane malalignment after tension-band plating removal²⁶. Funk et al. did not recommend hemiepiphysectomies in patients with late-onset tibial vara with a body mass index of >35 kg/m² or a mechanical axis deviation of >80 mm of varus²⁷. Hemiepiphysectomy should be considered in the treatment strategy for limb malalignment in patients with mucopolysaccharidosis Type IV-A²⁸. Kempainen et al. reported a high prevalence of incomplete follow-up (about 12%) in patients undergoing growth modulation, and, of these, nearly one-third required a surgical procedure other than implant removal²⁹.

Other Pediatric Conditions

In a systematic review of the prevalence of pin-track infection using external fixation, the pediatric population was shown to be at greatest risk³⁰. In a study on hexapod external fixators, Iobst et al. demonstrated that the universal joints attached to the undersurface of rings allowed more translational rotation, whereas the ball and socket joints attached to the outer surface of rings allowed more rotational correction and more correction with less strut change in patients with severe deformity³¹. The mechanisms by which isolated distal tibial rotation osteotomy corrects excessive tibial torsion in children with cerebral palsy were extensively investigated³². The results of a double-blind, multicenter, randomized controlled trial comparing Botox (botulinum toxin type A) injection with placebo in children undergoing lower-limb lengthening supported the use of Botox as an adjunct treatment in limb lengthening³³.

Adult Limb Reconstruction

Osteotomy About the Knee: High Tibial Osteotomy

Measurement of alignment before and after high tibial osteotomy was best performed with standing long-leg radiographs. The reproducibility and reliability of alignment measurement were better than for knee radiographs and were almost as good as a computed tomographic (CT) scan. The extra time and specialized software needed for CT scan measurement were not worthwhile³⁴. Single-photon emission CT and conventional CT (SPECT/CT) were used to evaluate knee compartments after opening-wedge high tibial osteotomy for providing clinical guidance for proper correction. Optimal realignment requires overcorrection that does not exceed 3° valgus. An increased signal was noted in the patellofemoral joint when the patellar position was changed^{35,36}.

A systematic review of 19 high tibial osteotomy studies including 1,189 patients showed that 87% returned to sport and 85% returned to work. Of those patients who did return to work and sport, 90% did so within 1 year. At a mean of 7 years, 93% of the patients did not progress to total knee replacement³⁷.

Medial opening-wedge high tibial osteotomy was performed on 18 patients with malunited tibial plateau fractures (excluding complex intra-articular deformity and severe arthritis) using a plate and bone graft. Coronal and sagittal plane alignment and the Knee injury and Osteoarthritis Outcome Score (KOOS) significantly improved with surgical procedures³⁸.

Osteotomy About the Knee: Distal Femoral Osteotomy

Distal femoral osteotomy using an opening-wedge technique on 41 patients with valgus deformity resulted in 95% accuracy of correction using mechanical axis deviation, lateral distal femoral angle, and medial proximal tibial angle measurements³⁹, and outcomes using the Short Form-36 (SF-36) and AAOS lower limb module (LLM) were significantly improved. In 10 knees with patellar subluxation, the patella congruence angle significantly improved after distal femoral osteotomy³⁹.

Opening-wedge distal femoral osteotomy was used in 29 knees, and the angular correction was 8°, changing the mechanical axis angle from 188° to 180°. Lower-limb length was not significantly changed. At 5 years, 91% did not progress to total knee replacement⁴⁰.

In patients with hypoplasia of the lateral femoral condyle, an intra-articular osteotomy with advancement of the lateral femoral condyle led to improvement in the tibiofemoral angle from 35° to 9° and satisfactory outcomes⁴¹.

Total Knee Arthroplasty

Outcomes at 1 year after total knee replacement with high tibial osteotomy were no different from those performed without previous high tibial osteotomy. Osteotomy is an effective means of delaying total knee replacement without compromise of outcome, especially when high tibial osteotomy techniques are opening-wedge with a plate or using external fixation⁴².

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

A systematic review of 8 studies examining total knee arthroplasty after medial opening and lateral closing-wedge high tibial osteotomy showed similar performance. Clinical and radiographic outcomes, including revision rates, did not differ; however, there were more surgical technical concerns in conversion to total knee arthroplasty in the lateral closing-wedge high tibial osteotomy group than in the medial opening-wedge high tibial osteotomy group⁴³.

Following total knee arthroplasty, functional outcomes of patients with >15 mm of postoperative limb-length discrepancy were lower than those with <15 mm of limb-length discrepancy. This calls attention to the importance of limb-length discrepancy as it relates to functional outcomes of primary total knee arthroplasty³⁵.

Alignment

Although 10% of patients perceive limb-length discrepancy after total knee replacement, this resolves in most within 3 months and is not correlated with mechanical alignment⁴⁴. A systematic review of 18 studies and 2,214 patients concluded that malalignment may correlate with lower patient-reported outcome scores. Hadi et al. called for larger longitudinal studies with standardized methods for assessing alignment⁴⁵.

Remote-Controlled Internal Lengthening Nail

A magnetic, remote-controlled intramedullary nail was used to successfully lengthen the femur and tibia in 23 patients. The mean lengthening was 48 mm, the mean angular correction was 16°, and the mean consolidation index was 1.12 months/cm. Although 8 patients had overlengthening, the nails were driven back to the desired length⁴⁶. Another study examined the experience with these nails in 10 patients with a mean limb-length discrepancy of 4.7 cm. In all patients, the limb-lengthening goals were reached within 0.5 cm after a mean time of 53 days. However, in 2 patients, mechanical failures with unintended shortening were observed. In 1 patient, nail breakage occurred⁴⁷. In another review of 9 patients, the mean lengthening (and standard deviation) was 34.7 ± 10.7 mm. All patients reached normal alignment and normal joint orientation. An unintentional loss of the achieved length during the consolidation phase was noticed in patients with delayed bone healing in 2 cases. In the first case (loss of 20-mm distraction), the nail could be re-distracted and the goal length was achieved. In the second case (loss of 10-mm distraction), the nail broke shortly after the diagnosis and the nail was exchanged. Weight-bearing restrictions are necessary until consolidation⁴⁸.

The use of blocking screws is necessary to correct and prevent deformity when using the internal lengthening nail technique. A systematic approach to the appropriate use of blocking screws in these deformities is described, including the reverse rule of thumb as a quick reference to determine the ideal location(s) and number of blocking screws⁴⁹.

A comparison of femoral lengthening with the internal nail and a monolateral fixator showed quicker osseous union

with the nail. The nailing also had fewer complications, was better tolerated by patients, and was perceived to have a better cosmetic result⁵⁰.

Strategies to avoid anterior nail impingement include the use of rigid straight reamers. In a series of 45 retrograde femoral lengthenings, the mean lengthening was 5.9 cm and the consolidation index was 1.05 months/cm. The mean distance from the osteotomy site to the intercondylar notch of the femur was 81 mm. The mean posterior cortex reaming thickness was 3.7 mm. No impingement-related complications or nail damage were observed⁵¹.

Many complications can be avoided with attention to technical detail and with increased experience. Technique papers for antegrade femoral lengthening and retrograde femoral lengthening using the magnetic internal nail include indications, technical tips, and pitfalls to avoid. Mechanical stability with optimal nail sizing, osteotomy location or technique, reaming technique, rate and rhythm of distraction, use of blocking screws, and adjuvant soft-tissue procedures are important to achieve success^{52,53}.

Other Adult Reconstruction

Neglected patellar tendon rupture with massive proximal patellar migration was successfully treated in 2 cases with patellar transport using a circular external fixator and staged allograft reconstruction⁵⁴. Neglected knee dislocation with residual deformity was successfully treated using a 2-stage strategy with gradual reduction using the Ilizarov technique and subsequent arthroscopic anterior and posterior cruciate ligament reconstruction⁵⁵. Knee flexion contractures were treated using a circular external fixator and gradual correction, and one-third of the patients were simultaneously treated for ankle equinus. The mean range of motion at the time of the final follow-up was -10° extension, 64° flexion, 9° ankle dorsiflexion, and 29° ankle plantar flexion. The arc of motion in the knee was unchanged but was increased in the ankle. The difference between preoperative and postoperative ranges of motion was significant. All but 2 patients were able to stand and walk with fewer aids or no aids⁵⁶.

Despite numerous studies that work to derive evidence-based recommendations for the prevention of pin-site infections, substantial controversy exists with regard to the optimal protocol. Kazmers et al. comprehensively evaluated the current literature to provide an overview of factors that may influence the incidence of pin-site infections in patients undergoing treatment with external fixators and provided a description of the preferred surgical and postoperative pin-site protocols that they employed⁵⁷.

Lengthening of a short residual tibia after transtibial amputation by >100% was achieved with no complications. Moreover, a special, custom-made, external fixator-prosthesis composite was described to allow for early weight-bearing and exercising⁵⁸. Both the Gigli saw and De Bastiani corticotomy techniques result in good bone formation following distraction

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

osteogenesis of the tibia. The anterior tibial cortex consolidates more slowly than the other cortices in both groups. This is likely due to deficient soft-tissue coverage and direct periosteal damage at the time of osteotomy⁵⁹.

Trauma

Successful reconstruction of bone defects remains a surgical challenge. A critical bone defect suggests a fracture gap that will not undergo healing without some form of intervention (e.g., bone-grafting). Currently, there is no consensus on the lower-limit size of a critical bone defect. Haines et al. studied 40 patients with open tibial diaphyseal fractures to define this lower limit⁶⁰. They included patients with gap sizes between 10 and 50 mm. Twenty-one patients (52.5%) developed a non-union. The authors concluded that a fracture gap after intramedullary nailing of <25 mm had a reasonable probability of achieving union, whereas a larger gap likely warrants secondary bone-grafting⁶⁰. Bone morphogenetic protein-2 (BMP-2) has been used as a bone graft in such defects. In a rat bone defect model, Glatt et al. demonstrated that, by optimizing the mechanical environment, one can enhance BMP-2's effect on bone growth⁶¹. They reported that either a medium-stiffness external fixator or reverse dynamization could decrease the amount of BMP-2 necessary to generate the same amount of bridging callus. Reverse dynamization was defined as beginning with a low-stiffness external fixator that changed to a stiff external fixator at 2 weeks after an osteotomy⁶¹.

Infected nonunions with associated bone defects remain a surgical challenge. The Ilizarov method of distraction osteogenesis with circular ring fixators has been shown to be successful in eradicating infection, equalizing limb lengths, and allowing soft-tissue lesions to heal⁶²⁻⁶⁵. This is especially true in the case of periarticular bone defects in which surface implants or intramedullary nails cannot achieve the stability required for successful reconstruction. Circular frames that are mounted juxta-articularly using 1.8-mm Ilizarov wires that are tensioned to 130 kg achieve the stability required to promote bone union, to correct deformity, and to eradicate infection. Eralp et al. retrospectively reported on 13 patients who underwent distal tibial periarticular reconstruction⁶⁶. The mean bone loss was 4.8 mm. At a mean follow-up of 36 months, 92% (12 patients) achieved union with a mean external fixation index of 29 days/cm⁶⁶. Repo et al. published a mean 6.6-year follow-up on 16 patients who underwent tibial reconstruction with a mean bone gain of 3.8 cm⁶⁷. All patients had use of a latissimus dorsi free flap for the soft-tissue defect. Bone transport was performed using various circular external fixators. The external fixation index was 54 days/cm. Although a lengthy and, at times, arduous process for the patient, the health-related quality-of-life measures that were reported were comparable with age-standardized population samples⁶⁷.

A hexapod circular fixator is extremely helpful in correcting complex deformities because the surgeon does not need to build a physical hinge at the apex of the deformity (as in a

classic Ilizarov circular frame). Instead, a virtual hinge is created through the computer software and an algorithm with an accompanying schedule dictates how to move the struts to make the bone straight. Hughes et al. used a temporary hexapod frame intraoperatively to acutely correct complex tibial deformities followed by the insertion of an intramedullary nail⁶⁸. Thirteen procedures were performed in 12 patients. The tibial mechanical axis was restored to anatomic values; however, 2 patients developed a common peroneal nerve palsy, and 1 patient developed a tibial artery pseudoaneurysm⁶⁸. Although the hexapod frame can correct complex deformities, its other benefit is the ability for gradual correction. This needs to be considered for large angular corrections.

Anatomic alignment in fracture reconstruction remains dogmatic. Weinberg et al. studied 37 fractured tibiae in cadaveric skeletons⁶⁹. Intra-articular fractures were excluded. They correlated knee arthritis with >5° of coronal malreduction ($p = 0.006$) and malrotation of >10° ($p = 0.004$)⁶⁹. Hip arthritis was associated with tibial shortening of >10 mm ($p = 0.009$). Avilucea et al. compared 266 patients with distal tibial fractures with respect to whether postoperative angulation of >5° occurred⁷⁰. The lateral distal tibial angle and anterior distal tibial angle were measured on postoperative tibial radiographs. Twenty-six percent of patients in the infrapatellar group compared with 3.8% of patients in the suprapatellar group had >5° of coronal or sagittal malalignment ($p < 0.0001$)⁷⁰.

Baruah published a case report on using a modified Ilizarov external fixator to gradually reduce a chronic patellar fracture nonunion⁷¹. Yang et al. reported on 7 patients with tibial plateau malunions who were treated with intra-articular osteotomies⁷². Preoperative planning was achieved with templating off a 3-dimensional printed model of the malunion.

Combining internal fixation (intramedullary nails or plates) with external fixation during limb lengthening and deformity correction is a strategy to decrease the amount of time that the patient needs to remain in the circular frame while the regenerate heals (latency phase). Burghardt et al. compared patients who underwent classic Ilizarov lengthening with a matched cohort who underwent lengthening over an intramedullary nail⁷³. The mean external fixation time for the group that underwent lengthening over an intramedullary nail was 2.6 months compared with 7.6 months in the classic Ilizarov lengthening group. Of note, there were 4 deep infections, increased blood loss, and an increase in surgical cost in the group who underwent lengthening over an intramedullary nail⁷³. Watanabe and Matsushita performed deformity correction of 21 femoral nonunions with a specific type of osteotomy at the nonunion site called the chipping technique⁷⁴. No bone grafts were utilized, and all nonunions healed with 1 procedure (19 cases) or 2 procedures (2 cases)⁷⁴. Azzam and El-Sayed performed lengthening over an existing femoral intramedullary nail for 13 diaphyseal femoral nonunions⁷⁵. The mean length achieved was 4 cm, and union occurred in 11 cases (85%)⁷⁵.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

Foot and Ankle**Ankle and Hindfoot**

Although supramalleolar osteotomy has been successful for treating distal tibial malalignment with ankle arthritis⁷⁶, neither supramalleolar osteotomy nor calcaneal osteotomy will correct an intra-articular varus or valgus deformity at its apex. Plafondplasty, an intra-articular realignment akin to hemiepiphysal elevation, has been implemented to remedy the asymmetric ankle joint line deformity at the source⁷⁷. The article by Al-Nammari and Myerson⁷⁷ was technique-based and highlighted the need for lateral ligament reconstruction for the joint line varus.

Forefoot

Ginés-Cespedosa et al. were unable to find a correlation between the presence of uncorrected hindfoot deformity and the outcome of hallux valgus corrective surgical procedures⁷⁸. This finding undermines the teaching that a valgus hindfoot creates hallux valgus.

Brachymetatarsia is amenable to distraction osteogenesis, with a low complication rate⁷⁹, but when the regenerate does not heal, the resulting defect can be treated with bone-grafting using a bioabsorbable hydroxyapatite-collagen composite⁸⁰. The authors emphasized the need for preservation of the periosteum and the existing thin regenerate bone during grafting. An alternative treatment for this defect could have been an injectable self-contained phosphate-releasing scaffold to promote biomineralization⁸¹.

Basic Science

Bhave et al. studied the effect of femoral lengthening, with either lengthening over a nail or Ilizarov classic methods, determining that patients with congenital femoral shortening who underwent lengthening had a higher incidence of loss of terminal knee flexion at a minimum follow-up of 2 years⁸². Burzyńska et al. showed that, in small children, these life-sized models (10 cm long) could be used to mount external fixators and to trial deformity correction surgical procedures⁸³. Morasiewicz et al. examined weight distribution and balance after Ilizarov osteotomy reconstruction surgical procedures

and found that limb realignment and length equalization restored these parameters to close to normal values⁸⁴.

Adjuvant Intervention for Bone Healing

Understanding the biological principles of distraction osteogenesis is the first step toward augmenting bone formation. A fantastic review focusing on similarities between clinically observed regenerate and basic in vivo tissue engineering was published⁸⁵. In a distraction osteogenesis model, researchers found that slower distraction rates were associated with osteogenesis originating from the bone end and a faster distraction rate created periosteal new bone with a thicker and more voluminous regenerate⁸⁶. In a meta-analysis of clinical distraction osteogenesis publications, both low-intensity pulsed ultrasound (LIPUS) and pulsed electromagnetic fields were shown to significantly reduce time to healing by a mean time of 11 days/cm⁸⁷. A separate meta-analysis of clinical distraction osteogenesis cases showed that LIPUS lowered healing time by a mean time of 15 days/cm⁸⁸.

Promising osteogenesis-promoting adjuvants include sclerostin antibody injection⁸⁹, BMP-2 with osteoprotegerin⁹⁰, human fetal mesenchymal stem cell secretome⁹¹, and staphylococcal enterotoxin⁹².

Reggie C. Hamdy, MD, MSc, FRCS(C)¹

Mitchell Bernstein, MD, FRCSC²

Austin T. Fragomen, MD³

S. Robert Rozbruch, MD³

¹Shriners Hospital for Children, Montreal, Quebec, Canada

²Loyola University Stritch School of Medicine, Maywood, Illinois

³Hospital for Special Surgery, New York, NY

E-mail address for R.C. Hamdy: rhamdy@shriners.mcgill.ca

ORCID iD for R.C. Hamdy: [0000-0002-0664-2843](https://orcid.org/0000-0002-0664-2843)

References

- Paley D. Why are deformity concepts still not a mainstream part of orthopaedics? *J Limb Lengthen Reconstr.* 2016;2:1-2.
- Hamdy RC. Evolution in long bone deformity correction in the post-Ilizarov era: external to internal devices. *J Limb Lengthen Reconstr.* 2016;2(2):61-7.
- Green SA. What happened to the regenerate? What would Ilizarov say? *J Limb Lengthen Reconstr.* 2016;2:3-5.
- Paley D. Surgical reconstruction for fibular hemimelia. *J Child Orthop.* 2016 Dec;10(6):557-83. Epub 2016 Dec 1.
- Paley D. Tibial hemimelia: new classification and reconstructive options. *J Child Orthop.* 2016 Dec;10(6):529-55. Epub 2016 Dec 1.
- Eidelman M, Jauregui JJ, Standard SC, Paley D, Herzenberg JE. Hip stability during lengthening in children with congenital femoral deficiency. *Int Orthop.* 2016 Dec;40(12):2619-25. Epub 2016 Sep 27.
- Saldanha KAN, Blakey CM, Broadley P, Fernandes JA. Defining patho-anatomy of the knee in congenital longitudinal lower limb deficiencies. *J Limb Lengthen Reconstr.* 2016;2:48-54.
- Shahcheraghi GH, Javid M. Functional assessment in tibial hemimelia (can we also save the foot in reconstruction?). *J Pediatr Orthop.* 2016 Sep;36(6):572-81.
- Borzunov DY, Chevardin AY, Mitrofanov AI. Management of congenital pseudarthrosis of the tibia with the Ilizarov method in a paediatric population: influence of aetiological factors. *Int Orthop.* 2016 Feb;40(2):331-9. Epub 2015 Nov 7.
- Alzahrani MM, Fassier F, Hamdy RC. Use of the Fassier-Duval telescopic rod for the management of congenital pseudarthrosis of the tibia. *J Limb Lengthen Reconstr.* 2016;2:23-8.
- Nomura I, Watanabe K, Matsubara H, Shirai T, Tsuchiya H. Plating following gradual realignment with the Taylor Spatial Frame for refractory congenital pseudarthrosis of the tibia: a novel technique. *J Limb Lengthen Reconstr.* 2016 (2):108-12.
- Iobst CA. Hexapod external fixation of tibia fractures in children. *J Pediatr Orthop.* 2016 Jun;36(Suppl 1):S24-8.
- Abdelkhalik M, El-Alfy B, Ali AM. Ilizarov bone transport versus fibular graft for reconstruction of tibial bone defects in children. *J Pediatr Orthop B.* 2016 Nov;25(6):556-60.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

- 14.** Shore BJ, DiMauro JP, Spence DD, Miller PE, Glotzbecker MP, Spencer S, Hedequist D. Uniplanar versus Taylor Spatial Frame external fixation for pediatric diaphyseal tibia fractures: a comparison of cost and complications. *J Pediatr Orthop.* 2016 Dec;36(8):821-8.
- 15.** Mills G, Nelson S. An improved spreadsheet for calculating limb length discrepancy and epiphysiodesis timing using the multiplier method. *J Child Orthop.* 2016 Aug;10(4):313-9. Epub 2016 Jun 29.
- 16.** Gaumétou E, Mallet C, Souchet P, Mazda K, Ilharreborde B. Poor efficiency of Eight-Plates in the treatment of lower limb discrepancy. *J Pediatr Orthop.* 2016 Oct-Nov;36(7):715-9.
- 17.** Makarov MR, Dunn SH, Singer DE, Rathjen KE, Ramo BA, Chukwuyerenwa CK, Birch JG. Complications associated with epiphysiodesis for management of leg length discrepancy. *J Pediatr Orthop.* 2016 Aug 19. [Epub ahead of print].
- 18.** Donnan LT, Gomes B, Donnan A, Harris C, Torode I, Heidt C. Ilizarov tibial lengthening in the skeletally immature patient. *Bone Joint J.* 2016 Sep;98-B(9):1276-82.
- 19.** Abousamra O, Orozco MDPD, Rogers KJ, lobst C, Nichols LR, Thacker M. Ankle alignment after tibial lengthening and syndesmosis fixation: a comparison study. *J Limb Lengthen Reconstr.* 2016;2:35-9.
- 20.** Yoshida Y, Matsubara H, Aikawa T, Ugaji S, Tsuchiya H. Complications related to fibula resection during tibial lengthening performed with the Taylor Spatial Frame. *J Limb Lengthen Reconstr.* 2016 (2):82-5.
- 21.** Chilbule SK, Dutt V, Madhuri V. Limb lengthening in achondroplasia. *Indian J Orthop.* 2016 Jul-Aug;50(4):397-405.
- 22.** Duque Orozco MD, Abousamra O, Chen BP, Rogers KJ, Sees JP, Miller F. Knee deformities in children with Down syndrome: a focus on knee malalignment. *J Pediatr Orthop.* 2016 Jun 18. [Epub ahead of print].
- 23.** Calis M, Oznur A, Ekin O, Vargel I. Correction of brachymetatarsia and medial angulation of the great toe of Apert foot by distraction osteogenesis: a review of 7 years of experience. *J Pediatr Orthop.* 2016 Sep;36(6):582-8.
- 24.** Brooks JT, Bernholt DL, Tran KV, Ain MC. The tibial slope in patients with achondroplasia: its characterization and possible role in genu recurvatum development. *J Pediatr Orthop.* 2016 Jun;36(4):349-54.
- 25.** Stevens PM. The role of guided growth as it relates to limb lengthening. *J Child Orthop.* 2016 Dec;10(6):479-86. Epub 2016 Dec 2.
- 26.** Farr S, Arabai HM, Meizer E, Ganger R, Radler C. Rebound of frontal plane malalignment after tension band plating. *J Pediatr Orthop.* 2016 Aug 29. [Epub ahead of print].
- 27.** Funk SS, Mignemi ME, Schoenecker JG, Lovejoy SA, Mencia GA, Martus JE. Hemiepiphysiodesis implants for late-onset tibia vara: a comparison of cost, surgical success, and implant failure. *J Pediatr Orthop.* 2016 Jan;36(1):29-35.
- 28.** Cooper GA, Southorn T, Eastwood DM, Bache CE. Lower extremity deformity management in MPS IVA, Morquio-Brailsford syndrome: preliminary report of hemiepiphysiodesis correction of genu valgum. *J Pediatr Orthop.* 2016 Jun;36(4):376-81.
- 29.** Kempainen JW, Hood KA, Roocroft JH, Schlechter JA, Edmonds EW. Incomplete follow-up after growth modulation surgery: incidence and associated complications. *J Pediatr Orthop.* 2016 Jul-Aug;36(5):516-20.
- 30.** lobst CA, Liu RW. A systematic review of incidence of pin track infections associated with external fixation. *J Limb Lengthen Reconstr.* 2016;2:6-16.
- 31.** lobst CA, Samchukov M, Cherkashin A. A comparison of deformity correction capabilities in hexapod frame systems. *J Limb Lengthen Reconstr.* 2016;2:29-34.
- 32.** Andrisevic E, Westberry DE, Pugh LI, Bagley AM, Tanner S, Davids JR. Correction of tibial torsion in children with cerebral palsy by isolated distal tibia rotation osteotomy: a short-term, in vivo anatomic study. *J Pediatr Orthop.* 2016 Oct-Nov;36(7):743-8.
- 33.** Hamdy RC, Montpetit K, Aiona MD, MacKenzie WG, van Bosse HJ, Narayanan U, Raney EM, Chafetz RS, Thomas SE, Weir S, Gregory S, Yorgova P, Takahashi S, Rinaldi M, Zhang X, Dahan-Oliel N. Safety and efficacy of botulinum toxin A in children undergoing lower limb lengthening and deformity correction: results of a double-blind, multicenter, randomized controlled trial. *J Pediatr Orthop.* 2016 Jan;36(1):48-55.
- 34.** Akamatsu Y, Sotozawa M, Kobayashi H, Kusayama Y, Kumagai K, Saito T. Usefulness of long tibial axis to measure medial tibial slope for opening wedge high tibial osteotomy. *Knee Surg Sports Traumatol Arthrosc.* 2016 Nov;24(11):3661-7. Epub 2014 Oct 29.
- 35.** Kim SH, Rhee SM, Lim JW, Lee HJ. The effect of leg length discrepancy on clinical outcome after TKA and identification of possible risk factors. *Knee Surg Sports Traumatol Arthrosc.* 2016 Aug;24(8):2678-85. Epub 2015 Nov 19.
- 36.** Kim TW, Kim BK, Kim DW, Sim JA, Lee BK, Lee YS. The SPECT/CT evaluation of compartmental changes after open wedge high tibial osteotomy. *Knee Surg Relat Res.* 2016 Dec;128(4):263-9.
- 37.** Ekhtiari S, Haldane CE, de Sa D, Simunovic N, Musahl V, Ayeni OR. Return to work and sport following high tibial osteotomy: a systematic review. *J Bone Joint Surg Am.* 2016 Sep 21;98(18):1568-77.
- 38.** Sundararajan SR, Nagaraja HS, Rajasekaran S. Medial open wedge high tibial osteotomy for varus malunited tibial plateau fractures. *Arthroscopy.* 2017 Mar;33(3):586-94. Epub 2016 Nov 19.
- 39.** Elattar O, Swarup I, Lam A, Nguyen J, Fragomen A, Rozbruch SR. Open wedge distal femoral osteotomy: accuracy of correction and patient outcomes. *HSS J.* 2016 Jul;1:8.
- 40.** Madelaine A, Lording T, Villa V, Lustig S, Servien E, Neyret P. The effect of lateral opening wedge distal femoral osteotomy on leg length. *Knee Surg Sports Traumatol Arthrosc.* 2016 Mar;24(3):847-54. Epub 2014 Oct 19.
- 41.** Feldman DS, Goldstein RY, Kurland AM, Sheikh Taha AM. Intra-articular osteotomy for genu valgum in the knee with a lateral compartment deficiency. *J Bone Joint Surg Am.* 2016 Jan 20;98(2):100-7.
- 42.** W-Dahl A, Robertsson O. Similar outcome for total knee arthroplasty after previous high tibial osteotomy and for total knee arthroplasty as the first measure. *Acta Orthop.* 2016 Aug;87(4):395-400. Epub 2016 Jun 24.
- 43.** Han JH, Yang JH, Bhandare NN, Suh DW, Lee JS, Chang YS, Yeom JW, Nha KW. Total knee arthroplasty after failed high tibial osteotomy: a systematic review of open versus closed wedge osteotomy. *Knee Surg Sports Traumatol Arthrosc.* 2016 Aug;24(8):2567-77. Epub 2015 Sep 30.
- 44.** Goldstein ZH, Yi PH, Batko B, Kearns S, Tetreault MW, Levine BR, Della Valle CJ, Sporer SM. Perceived leg-length discrepancy after primary total knee arthroplasty: does knee alignment play a role? *Am J Orthop (Belle Mead NJ).* 2016 Nov/Dec;45(7):E429-33.
- 45.** Hadi M, Barlow T, Ahmed I, Dunbar M, McCulloch P, Griffin D. Does malalignment affect patient reported outcomes following total knee arthroplasty: a systematic review of the literature. *Springerplus.* 2016 Jul 28;5(1):1201.
- 46.** Karakoyun O, Sokucu S, Erol MF, Kucukkaya M, Kabukcuoglu YS. Use of a magnetic bone nail for lengthening of the femur and tibia. *J Orthop Surg (Hong Kong).* 2016 Dec;24(3):374-8.
- 47.** Tiefenboeck TM, Zak L, Bukaty A, Wozasek GE. Pitfalls in automatic limb lengthening - first results with an intramedullary lengthening device. *Orthop Traumatol Surg Res.* 2016 Nov;102(7):851-5. Epub 2016 Aug 12.
- 48.** Wiebking U, Liodakis E, Kenaway M, Krettek C. Limb lengthening using the PRECICE(TM) Nail System: complications and results. *Arch Trauma Res.* 2016 Aug 23;5(4):e36273.
- 49.** Muthusamy S, Rozbruch SR, Fragomen AT. The use of blocking screws with internal lengthening nail and reverse rule of thumb for blocking screws in limb lengthening and deformity correction surgery. *Strategies Trauma Limb Reconstr.* 2016 Nov;11(3):199-205. Epub 2016 Sep 24.
- 50.** Laubscher M, Mitchell C, Timms A, Goodier D, Calder P. Outcomes following femoral lengthening: an initial comparison of the Precice intramedullary lengthening nail and the LRS external fixator monorail system. *Bone Joint J.* 2016 Oct;98-B(10):1382-8.
- 51.** Kucukkaya M, Karakoyun Ö, Erol MF. The importance of reaming the posterior femoral cortex before inserting lengthening nails and calculation of the amount of reaming. *J Orthop Surg Res.* 2016 Jan 16;11:11.
- 52.** Fragomen AT, Rozbruch SR. Lengthening of the femur with a remote-controlled magnetic intramedullary nail: retrograde technique. *JBJS Essent Surg Tech.* 2016 May 11;6(2):e20.
- 53.** Rozbruch SR, Fragomen AT. Lengthening of the femur with a remote-controlled magnetic intramedullary nail: anterograde technique. *JBJS Essent Surg Tech.* 2016 Jan 13;6(1):e2.
- 54.** Elattar O, Coleman SH, Warren RF, Rozbruch SR. Neglected patellar tendon rupture with massive proximal patellar migration treated with patellar transport and staged allograft reconstruction: a report of 2 cases. *Orthop J Sports Med.* 2016 Nov 21;4(11):2325967116672175.
- 55.** Polyzois VD, Stathopoulos IP, Benetos IS, Pneumaticsos SG. A two-stage procedure for the treatment of a neglected posterolateral knee dislocation: gradual reduction with an Ilizarov external fixator followed by arthroscopic anterior and posterior cruciate ligament reconstruction. *Knee.* 2016 Jan;23(1):181-4. Epub 2015 Dec 10.
- 56.** Vulcano E, Markowitz JS, Fragomen AT, Rozbruch SR. Gradual correction of knee flexion contracture using external fixation. *J Limb Lengthen Reconstr.* 2016;2:102-7.
- 57.** Kazmers NH, Fragomen AT, Rozbruch SR. Prevention of pin-site infection in external fixation: a review of the literature. *Strateg Trauma Limb Reconstr.* 2016;11(2):75-85.
- 58.** Lam A, Garrison G, Rozbruch SR. Lengthening of tibia after trans-tibial amputation: use of a weight bearing external fixator-prosthesis composite. *HSS J.* 2016 Feb;12(1):85-90. Epub 2015 Sep 8.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

- 59.** Peek AC, Timms A, Chin KF, Calder P, Goodier D. Patterns of healing: a comparison of two proximal tibial osteotomy techniques. *Strategies Trauma Limb Reconstr.* 2016 Apr;11(1):59-62. Epub 2016 Feb 16.
- 60.** Haines NM, Lack WD, Seymour RB, Bosse MJ. Defining the lower limit of a "critical bone defect" in open diaphyseal tibial fractures. *J Orthop Trauma.* 2016 May;30(5):e158-63.
- 61.** Glatt V, Bartnikowski N, Quirk N, Schuetz M, Evans C. Reverse dynamization: influence of fixator stiffness on the mode and efficiency of large-bone-defect healing at different doses of rhBMP-2. *J Bone Joint Surg Am.* 2016 Apr 20;98(8):677-87.
- 62.** Rohilla R, Siwach K, Devgan A, Singh R, Wadhvani J, Ahmed N. Outcome of distraction osteogenesis by ring fixator in infected, large bone defects of tibia. *J Clin Orthop Trauma.* 2016 Oct-Dec;7(Suppl 2):201-9. Epub 2016 Apr 20.
- 63.** Rohilla R, Wadhvani J, Devgan A, Singh R, Khanna M. Prospective randomised comparison of ring versus nail fixator in infected gap nonunion of tibia treated with distraction osteogenesis. *Bone Joint J.* 2016 Oct;98-B(10):1399-405.
- 64.** Sadek AF, Laklok MA, Foully EH, Elshafie M. Two stage reconstruction versus bone transport in management of resistant infected tibial diaphyseal nonunion with a gap. *Arch Orthop Trauma Surg.* 2016 Sep;136(9):1233-41. Epub 2016 Jul 22.
- 65.** Semaya AelS, Badawy E, Hasan M, El-Nakeeb RM. Management of post-traumatic bone defects of the tibia using vascularised fibular graft combined with Ilizarov external fixator. *Injury.* 2016 Apr;47(4):969-75. Epub 2016 Feb 11.
- 66.** Eralp IL, Kocaoğlu M, Dikmen G, Azam ME, Balcı HI, Bilen FE. Treatment of infected nonunion of the juxta-articular region of the distal tibia. *Acta Orthop Traumatol Turc.* 2016;50(2):139-46.
- 67.** Repo JP, Barner-Rasmussen I, Roine RP, Sintonen H, Tukiainen EJ. Treatment of compound tibia fracture with microvascular latissimus dorsi flap and the Ilizarov technique: a cross-sectional study of long-term outcomes. *J Plast Reconstr Aesthet Surg.* 2016 Apr;69(4):524-32. Epub 2016 Jan 14.
- 68.** Hughes A, Parry M, Heidari N, Jackson M, Atkins R, Monsell F. Computer hexapod-assisted orthopaedic surgery for the correction of tibial deformities. *J Orthop Trauma.* 2016 Jul;30(7):e256-61.
- 69.** Weinberg DS, Park PJ, Liu RW. Association between tibial malunion deformity parameters and degenerative hip and knee disease. *J Orthop Trauma.* 2016 Sep;30(9):510-5.
- 70.** Avilucea FR, Triantafyllou K, Whiting PS, Perez EA, Mir HR. Suprapatellar intramedullary nail technique lowers rate of malalignment of distal tibia fractures. *J Orthop Trauma.* 2016 Oct;30(10):557-60.
- 71.** Baruah RK. Modified Ilizarov in difficult fracture of the patella. A case report. *J Orthop Case Rep.* 2016 Jan-Mar;6(1):26-8.
- 72.** Yang P, Du D, Zhou Z, Lu N, Fu Q, Ma J, Zhao L, Chen A. 3D printing-assisted osteotomy treatment for the malunion of lateral tibial plateau fracture. *Injury.* 2016 Dec;47(12):2816-21. Epub 2016 Sep 12.
- 73.** Burghardt RD, Manzotti A, Bhavé A, Paley D, Herzenberg JE. Tibial lengthening over intramedullary nails: a matched case comparison with Ilizarov tibial lengthening. *Bone Joint Res.* 2016 Jan;5(1):1-10.
- 74.** Watanabe Y, Matsushita T. Femoral non-union with malalignment: reconstruction and biological stimulation with the chipping technique. *Injury.* 2016 Dec;47(Suppl 6):S47-52.
- 75.** Azzam W, El-Sayed M. Ilizarov distraction osteogenesis over the preexisting nail for treatment of nonunited femurs with significant shortening. *Eur J Orthop Surg Traumatol.* 2016 Apr;26(3):319-28. Epub 2016 Jan 25.
- 76.** Krause F, Vejjkovic A, Schmid T. Supramalleolar osteotomies for posttraumatic malalignment of the distal tibia. *Foot Ankle Clin.* 2016 Mar;21(1):1-14.
- 77.** Al-Nammari SS, Myerson MS. The use of tibial osteotomy (ankle plafondplasty) for joint preservation of ankle deformity and early arthritis. *Foot Ankle Clin.* 2016 Mar;21(1):15-26.
- 78.** Ginés-Cespedosa A, Pérez-Prieto D, Muñetón D, González-Lucena G, Millán A, de Zabala S, Busquets R. Influence of hindfoot malalignment on hallux valgus operative outcomes. *Foot Ankle Int.* 2016 Aug;37(8):842-7. Epub 2016 Apr 20.
- 79.** Hosny GA, Ahmed AS. Distraction osteogenesis of fourth brachymetatarsia. *Foot Ankle Surg.* 2016 Mar;22(1):12-6. Epub 2015 Apr 8.
- 80.** Hamada M, Sakamoto Y, Nagasao T, Kishi K. Treatment of complications after distraction osteogenesis for brachymetatarsia of the fourth metatarsal. *Plast Reconstr Surg Glob Open.* 2016 Jul 21;4(7):e817.
- 81.** Nayef L, Mekhail M, Benameur L, Rendon JS, Hamdy R, Tabrizian M. A combinatorial approach towards achieving an injectable, self-contained, phosphate-releasing scaffold for promoting biomineralization in critical size bone defects. *Acta Biomater.* 2016 Jan;29:389-97. Epub 2015 Oct 17.
- 82.** Bhavé A, Shabtai L, Woelber E, Apelyan A, Paley D, Herzenberg JE. Muscle strength and knee range of motion after femoral lengthening. *Acta Orthop.* 2017 Apr;88(2):179-84. Epub 2016 Nov 28.
- 83.** Burzyńska K, Morasiewicz P, Filipiak J. The use of 3D printing technology in the Ilizarov method treatment: pilot study. *Adv Clin Exp Med.* 2016 Nov-Dec;25(6):1157-63.
- 84.** Morasiewicz P, Dragan S, Dragan SL, Wrzosek Z, Pawik L. Pedobarographic analysis of body weight distribution on the lower limbs and balance after Ilizarov corticotomies. *Clin Biomech (Bristol, Avon).* 2016 Jan;31:2-6. Epub 2015 Oct 23.
- 85.** Dhaliwal K, Kunchur R, Farhadieh R. Review of the cellular and biological principles of distraction osteogenesis: an in vivo bioreactor tissue engineering model. *J Plast Reconstr Aesthet Surg.* 2016 Feb;69(2):e19-26. Epub 2015 Nov 25.
- 86.** Saulacic N, Nakahara K, Iizuka T, Haga-Tsujimura M, Hofstetter W, Scolozzi P. Comparison of two protocols of periosteal distraction osteogenesis in a rabbit calvaria model. *J Biomed Mater Res B Appl Biomater.* 2016 Aug;104(6):1121-31. Epub 2015 Jun 2.
- 87.** Jauregui JJ, Ventimiglia AV, Grieco PW, Frumberg DB, Herzenberg JE. Regenerate bone stimulation following limb lengthening: a meta-analysis. *BMC Musculoskelet Disord.* 2016 Sep 29;17(1):407-14.
- 88.** Raza H, Saltaji H, Kaur H, Flores-Mir C, El-Bialy T. Effect of low-intensity pulsed ultrasound on distraction osteogenesis treatment time: a meta-analysis of randomized clinical trials. *J Ultrasound Med.* 2016 Feb;35(2):349-58. Epub 2016 Jan 18.
- 89.** Alzahrani MM, Rauch F, Hamdy RC. Does sclerostin depletion stimulate fracture healing in a mouse model? *Clin Orthop Relat Res.* 2016 May;474(5):1294-302. Epub 2015 Nov 25.
- 90.** Bougioukli S, Jain A, Sugiyama O, Tinsley BA, Tang AH, Tan MH, Adams DJ, Kostenuik PJ, Lieberman JR. Combination therapy with BMP-2 and a systemic RANKL inhibitor enhances bone healing in a mouse critical-sized femoral defect. *Bone.* 2016 Mar;84:93-103. Epub 2015 Dec 23.
- 91.** Xu J, Wang B, Sun Y, Wu T, Liu Y, Zhang J, Lee WY, Pan X, Chai Y, Li G. Human fetal mesenchymal stem cell secretome enhances bone consolidation in distraction osteogenesis. *Stem Cell Res Ther.* 2016 Sep 10;7(1):134-45.
- 92.** Xu J, Wu T, Sun Y, Wang B, Zhang J, Lee WY, Chai Y, Li G. Staphylococcal enterotoxin C2 expedites bone consolidation in distraction osteogenesis. *J Orthop Res.* 2016 Jul 19:1-10. Epub 2016 Jul 19.